



## Product Information Sheet

---

### Human IL-15 ELISA Kit

<b>Catalog No.</b>	EK0426
<b>Size</b>	96T
<b>Range</b>	15.6pg/ml-1000pg/ml
<b>Sensitivity</b>	< 3 pg/ml

#### **Specificity**

No detectable cross-reactivity with any other cytokine.

#### **Storage**

Store at 4°C for frequent use, at -20°C for infrequent use.

Avoid multiple freeze-thaw cycles (Shipped with wet ice.)

#### **Expiration**

Four months at 4°C and eight months at -20°C.

#### **Application**

For quantitative detection of human IL-15 in sera, plasma, body fluids, tissue lysates or cell culture supernates.

#### **Principle**

Human IL-15 ELISA Kit was based on standard sandwich enzyme-linked immune-sorbent assay technology. Human IL-15 specific polyclonal antibodies were precoated onto 96-well plates. The human specific detection polyclonal antibodies were biotinylated. The test samples and biotinylated detection antibodies were added to the wells subsequently and then followed by washing with PBS or TBS buffer. Avidin-Biotin-Peroxidase Complex was added and unbound conjugates were washed away with PBS or TBS buffer. HRP substrate TMB was used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the human IL-15 amount of sample captured in plate.

#### **Kit Components**

1. Lyophilized recombinant human IL-15 standard: 10ng/tubex2.
2. One 96-well plate precoated with anti- human IL-15 antibody.
3. Sample diluent buffer: 30 ml
4. Biotinylated anti- human IL-15 antibody : 130µl, dilution 1:100.
5. Antibody diluent buffer: 12ml.
6. Avidin-Biotin-Peroxidase Complex (ABC) : 130µl, dilution 1:100.
7. ABC diluent buffer: 12ml.
8. TMB color developing agent: 10ml.
9. TMB stop solution: 10ml.

#### **Material Required But Not Provided**

1. Microplate reader in standard size. and Automated plate washer.
2. Adjustable pipettes and pipette tips. Multichannel pipettes are recommended in the condition of large amount of samples in the detection.
3. Clean tubes and Eppendorf tubes.
4. Washing buffer (neutral PBS or TBS).

Preparation of 0.01M **TBS**: Add 1.2g Tris, 8.5g NaCl; 450µl of purified acetic acid or 700µl of concentrated hydrochloric acid to 1000ml H<sub>2</sub>O and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

Preparation of 0.01 M **PBS**: Add 8.5g sodium chloride, 1.4g Na<sub>2</sub>HPO<sub>4</sub> and 0.2g NaH<sub>2</sub>PO<sub>4</sub> to 1000ml distilled water and adjust pH to 7.2-7.6. Finally, adjust the total volume to 1L.

**To reorder contact us at:**  
**Antagene, Inc.**  
**Toll Free: 1(866)964-2589**  
**Tel: (650) 964-2589**  
**Fax: (650) 964-2519**  
**email: Info@antageneinc.com**

**FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC AND CLINICAL USE.**


# Product Information Sheet

---

## **Notice for Application of Kit**

1. Before using Kit, spin tubes and bring down all components to bottom of tube.
2. Duplicate well assay was recommended for both standard and sample testing.
3. Don't let 96-well plate dry, dry plate will inactivate active components on plate.
4. In order to avoid marginal effect of plate incubation due to temperature difference ( reaction may be stronger in the marginal wells), it is suggested that the diluted ABC and TMB solution will be pre-warmed in 37°C for 30 min before using.

## **Human IL-15 ELISA Kit-1X96 Well Plate Image**



## **Background**

Interleukin (IL)-15 is a cytokine with the ability to stimulate the proliferation activity of Th1 and/or Th2 lymphocytes. IL-15 is a novel cytokine whose effects on T-cell activation and proliferation are similar to those of interleukin-2 (IL-2), presumably because IL-15 utilizes the beta and gamma chains of the IL-2 receptor.<sup>1</sup> IL-15 can play a role in the initiation and outcome of acute and chronic rejection. Anti-IL-15 therapy in combination with classic immunosuppression therapy might thus be beneficial in the prevention of acute, and especially chronic, allograft rejection.<sup>2</sup> The human IL15 gene is mapped to human chromosome 4q31 by fluorescence in situ hybridization.<sup>1</sup> The standard product used in this kit is recombinant human IL-15, consisting of 114 amino acids with the molecular mass of 12.9KDa.

## **Reference**

1. Anderson, D. M.; Johnson, L.; Glaccum, M. B.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Valentine, V.; Kirstein, M. N.; Shapiro, D. N.; Morris, S. W.; Grabstein, K.; Cosman, D. Chromosomal assignment and genomic structure of IL15. *Genomics* 25: 701-706, 1995.
2. Conti F, Frappier J, Dharancy S, Chereau C, Houssin D, Weill B, Calmus Y. Interleukin-15 production during liver allograft rejection in humans. *Transplantation*. 2003 Jul 15;76(1):210-6.