Cat. #: 60B307

Description:

The HIF1AN (Hypoxia-inducible factor 1 alpha inhibitor)hydroxylates HIF-1 alpha at Asp-803 in the C-terminal transactivation domain (CAD). It functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents interaction of HIF-1 with transcriptional coactivators including Cbp/p300-interacting transactivator. The HIF1AN is involved in transcriptional repression through interaction with HIF1A, VHL and histone deacetylases.

Immunogen/Specificity:

Polyclonal antibody produced in rabbits immunizing with a synthetic peptide corresponding to Middle residues of human HIF1AN (Hypoxia-inducible factor 1 alpha inhibitor)

References

Mahon,P.C., et al, Genes Dev. 15 (20), 2675-2686 (2001) Lando,D., et al, Genes Dev. 16 (12), 1466-1471 (2002) Hewitson,K.S., et al, J. Biol. Chem. 277 (29), 26351-26355 (2002) Elkins,J.M., et al, J. Biol. Chem. 278 (3), 1802-1806 (2003) Dann,C.E. III, et al, Proc. Natl. Acad. Sci. U.S.A. 99 (24), 15351-15356 (2002) Clone Number: Isotype: Species: Human, Mouse Storage and Stability: at -20oC

Storage buffer: This antibody is stored in PBS, 0.01% sodium azide and 50% glycerol.

Preparation: Purified by antigen-specific affinity chromatography.

Applications : ELISA Western Blotting (1µg/ml for 2hrs)