



Product Information Sheet

Polyclonal Anti-Mitogen-activated protein organizer 1, *MORG1* (Magnetic Bead Conjugate)

Catalogue No. PA1053-M

Lot No. 03A10

Ig type: rabbit IgG

Size: 100µg/vial

Specificity

Human, mouse, rat.

No cross reactivity with other proteins.

Recommended application

ImmunoPrecipitation (IP)

Immunogen

A synthetic peptide corresponding to a sequence near the N-terminal of human MORG1, identical to the related mouse and rat sequence.

Purity

Immunogen affinity purified.

Contents

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN₃.

Storage

Store at 4°C for frequent use.

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified magnetic beads. It is useful for immunoprecipitation

BACKGROUND

MORG1 (mitogen-activated protein kinase organizer 1), a member of the WD-40 protein family that was isolated as a binding partner of the extracellular signal-regulated kinase (ERK) pathway scaffold protein MP1. MORG1 specifically associates with several components of the ERK pathway, including MP1, Raf-1, MEK, and ERK, and stabilizes their assembly into an oligomeric complex. MORG1 facilitates ERK activation when cells are stimulated with lysophosphatidic acid, phorbol 12-myristate 13-acetate, or serum, but not in response to epidermal growth factor. Suppression of MORG1 by short interfering RNA leads to a marked reduction in ERK activity when cells are stimulated with serum. MORG1 is a component of a modular scaffold system that participates in the regulation of agonist-specific ERK signaling.

REFERENCE

Modular construction of a signaling scaffold: MORG1 interacts with components of the ERK cascade and links ERK signaling to specific agonists. Vomastek T, Schaeffer HJ, Tarcsfalvi A, Smolkin ME, Bissonette EA, Weber MJ. Department of Microbiology and Cancer Center, University of Virginia, Charlottesville, VA 22908, USA.