



Product Information Sheet

Polyclonal Anti-MMP16 (Magnetic Bead Conjugate)

Catalogue No. PA1123-M	Immunogen A synthetic peptide mapping at the C-terminal of human MMP16, identical to the related rat sequence.
Lot No. 08J01	
Ig type: rabbit IgG1	Purification Immunogen affinity purified
Size: 100µg/Vial	Contents Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN ₃ .
Specificity Human, rat, mouse. No cross reactivity with other proteins.	Storage Store at 4°C for frequent use.
Recommended application <i>Immunoprecipitation(IP)</i>	Description: This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified magnetic beads. It is useful for immunoprecipitation

BACKGROUND

The matrix metalloproteinase 16(MMP16) protein consists of 604 amino acids and has a characteristic MMP domain structure, which gene is mapped on human chromosome 8q21¹. Additionally, MMP16 has a C-terminal extension containing a potential transmembrane domain, similar to MMP14, MMP15, and MMP17². Furthermore, it is membrane-bound and is a member of the membrane-type MMPs that are a subclass in the MMP family since the other members lack a C-terminal transmembrane domain and are secreted as soluble forms². MMP16 is expressed as a 12-kb transcript in brain, placenta, heart, and some carcinoma cell lines, but is not detectably expressed in lung, kidney, liver, spleen, and muscle².

REFERENCE

- 1.Sato, H.; Tanaka, M.; Takino, T.; Inoue, M.; Seiki, M. : Assignment of the human genes for membrane-type-1, -2, and -3 matrix metalloproteinases (MMP14, MMP15, and MMP16) to 14q12.2, 16q12.2-q21, and 8q21, respectively, by in situ hybridization. *Genomics* 39: 412-413, 1997.
- 2.Takino, T.; Sato, H.; Shinagawa, A.; Seiki, M. : Identification of the second membrane-type matrix metalloproteinase (MT-MMP-2) gene from a human placenta cDNA library: MT-MMPs form a unique membrane-type subclass in the MMP family. *J. Biol. Chem.* 270: 23013-23020, 1995.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com