



Polyclonal Anti- Interleukin-1 alpha, *IL-1α* (Sepharose Bead Conjugate)

Catalogue No. PA1375-S

Lot No. 0131112067527

Ig type: rabbit IgG

Size: 100µg/vial

Specificity

Mouse. No cross reactivity with other proteins.

Recommended application

(Immunoprecipitation(IP))

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminal of mouse IL-1α (256-270 aa), identical to the related mouse and rat sequence.

Purification

Immunogen affinity purified.

Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Storage

Store at 4°C for frequent use.

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays

BACKGROUND

Interleukin-1 alpha (IL-1α) is a protein that in humans is encoded by the IL1A gene. The protein encoded by this gene is a cytokine of the interleukin-1 family. Interleukin-1 alpha possesses a wide spectrum of metabolic, physiological, haematopoietic activities, and plays one of the central roles in the regulation of the immune responses. It binds to the interleukin-1 receptor. Silver et al. (1990) showed that the mouse Il1a and Il1b genes are contained in a genomic fragment of about 70 kb. Further studies suggested that Il1b lies 5-prime to Il1a, that the 2 genes are oriented in the same direction, and that they are separated by about 50 kb.

REFERENCE

- 1.Nicklin MJ, Weith A, Duff GW (Jun 1994). "A physical map of the region encompassing the human interleukin-1 alpha, interleukin-1 beta, and interleukin-1 receptor antagonist genes". *Genomics* 19 (2): 382–4.
- 2.March CJ, Mosley B, Larsen A, Cerretti DP, Braedt G, Price V, Gillis S, Henney CS, Kronheim SR, Grabstein K, et al. (Aug 1985). "Cloning, sequence and expression of two distinct human interleukin-1 complementary DNAs". *Nature* 315 (6021): 641–7.
- 3.Bankers-Fulbright JL, Kalli KR, McKean DJ (1996). "Interleukin-1 signal transduction". *Life Sci.* 59 (2): 61–83.
- 4.Dinarelli CA (June 1997). "Induction of interleukin-1 and interleukin-1 receptor antagonist". *Semin. Oncol.* 24 (3 Suppl 9): S9–81–S9–93.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com