



---

## Product Information Sheet

---

### Polyclonal Anti- Synaptopodin (Magnetic Bead Conjugate)

**Catalogue No.** PA1384-M

**Immunogen**

A synthetic peptide corresponding to a sequence at the middle region of human synaptopodin (495-509 aa), identical to the related mouse and rat sequence.

**Lot No.** 0131112028427

**Ig type** rabbit IgG

**Purity**

Immunogen affinity purified.

**Size** 100µg/vial

**Contents**

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN<sub>3</sub>.

**Specificity**

Human, rat.

No cross reactivity with other proteins.

**Storage**

Store at 4°C for frequent use.

**Recommended application**

immunoprecipitation.

**Description**

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified magnetic beads. It is useful for immunoprecipitation.

#### BACKGROUND

The spine apparatus (SA) is a specialized form of endoplasmic reticulum (ER) that is found in a subpopulation of dendritic spines in central neurons. The SA consists of a series of stacked discs that are thought to be connected to each other and to the dendritic system of ER-tubules. The actin binding protein synaptopodin (which has originally been described in podocytes of the kidney) is an essential component of the SA. Mice that lack the gene for synaptopodin do not form a spine apparatus. The SA is believed to play a critical role in learning and memory. In summary, an important function of the spine apparatus is the regulation of plasticity at individual synapses, a process known as metaplasticity. The International Radiation Hybrid Mapping Consortium mapped the SYNPO gene to chromosome 5.

#### REFERENCE

- 1.Gray (1959). "Electron microscopy of synaptic contacts on dendrite spines of the cerebral cortex". Nature 183 (4675): 1592–3.
- 2.Cooney; Hurlburt, JL; Selig, DK; Harris, KM; Fiala, JC (2002). "Endosomal compartments serve multiple hippocampal dendritic spines from a widespread rather than a local store of recycling membrane". The Journal of neuroscience : the official journal of the Society for Neuroscience 22 (6): 2215–24.
- 3.Deller; Merten, T; Roth, SU; Mundel, P; Frotscher, M (2000). "Actin-associated protein synaptopodin in the rat hippocampal formation: localization in the spine neck and close association with the spine apparatus of principal neurons". The Journal of Comparative Neurology 418 (2): 164–81

**For Research Use Only not for diagnostic and clinical use**

**Contact:** Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: [Info@antageneinc.com](mailto:Info@antageneinc.com)