



## Mouse Monoclonal Antibody **NLRP3** conjugated to Sepharose Beads

CatalogNo: **ANT8024-S**

Size 200ul

Storage Store at 4 °C for frequent use

### Description

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays.

### **NLRP3 (ANT0049R) Rabbit mAb**

Formulation: 50% slurry in PBS pH 7.2 with 0.01mg NaN<sub>3</sub> preservative.

#### Host Species

- Rabbit
- Human, Mouse, Rat,

#### Reactivity

- WB, IHC, IF, IP, ELISA

#### Applications

#### MW

- 115kD (Observed)
- IgG, Kappa

#### Isotype

## Recommended Dilution Ratios

### IP

### Basic Information

**Clonality** Monoclonal

**Clone Number** ANT0049R


### Immunogen Information

**Specificity** Endogenous


### Target Information

Gene name	NLRP3		
Protein Name	NACHT LRR and PYD domains-containing protein 3		
	Organism	Gene ID	UniProt ID
	Human	<a href="#">114548</a> ;	<a href="#">Q96P20</a> ;
	Mouse		<a href="#">Q8R4B8</a> ;
Cellular Localization	Cytoplasm, Nuclear		
Function	<p>Disease:Defects in NLRP3 are a cause of Muckle-Wells syndrome (MWS) [MIM:191900]; also known as urticaria-deafness-amyloidosis syndrome. MWS is a hereditary periodic fever syndrome characterized by fever, chronic recurrent urticaria, arthralgias, progressive sensorineural deafness, and reactive renal amyloidosis. The disease may be severe if generalized amyloidosis occurs.,Disease:Defects in NLRP3 are the cause of chronic infantile neurologic cutaneous and articular syndrome (CINCA) [MIM:607115]; also known as 'neonatal onset multisystem inflammatory disease,' or NOMID, a rare congenital inflammatory disorder characterized by a triad of neonatal onset of cutaneous symptoms, chronic meningitis, and joint manifestations with recurrent fever and inflammation.,Disease:Defects in NLRP3 are the cause of familial cold autoinflammatory syndrome type 1 (FCAS1) [MIM:120100]; commonly known as familial cold urticaria. FCAS are rare autosomal dominant systemic inflammatory diseases characterized by episodes of rash, arthralgia, fever and conjunctivitis after generalized exposure to cold.,Function:May function as an inducer of apoptosis. Interacts selectively with ASC and this complex may function as an upstream activator of NF-kappa-B signaling. Inhibits TNF-alpha induced activation and nuclear translocation of RELA/NF-KB p65. Also inhibits transcriptional activity of RELA. Activates caspase-1 in response to a number of triggers including bacterial or viral infection which leads to processing and release of IL1B and IL18.,induction:By TNFalpha.,online information:Repertory of FMF and hereditary autoinflammatory disorders mutations,similarity:Belongs to the NLRP family.,similarity:Contains 1 DAPIN domain.,similarity:Contains 1 NACHT domain.,similarity:Contains 7 LRR (leucine-rich) repeats.,subunit:Interacts with PYCARD/ASC. Part of the NALP3 inflammasome complex which is involved in activation of caspase-1 and caspase-5, leading to processing of IL1B and IL18.,tissue specificity:Expressed in blood leukocytes. Strongly expressed in polymorphonuclear cells and osteoblasts. Undetectable or expressed at a lower magnitude in B- and T-lymphoblasts, respectively. High level of expression detected in chondrocytes. Detected in non-keratinizing epithelia of oropharynx, esophagus and ectocervix and in the urothelial layer of the bladder.,</p>		


## Validation Data




Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-NLRP3 (ANT0049R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: J774A.1 Lane 2: RAW364.7 Lane 3: THP-1 Predicted band size: 118kDa Observed band size: 118kDa




Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100



Western blot analysis of lysate from K562 cells, using NLRP3 Antibody.



Rat kidney tissue was stained with Anti-NLRP3 (ANT0049R) rabbit Antibody

For Research use only, not for diagnostics and clinical use  
Contact Antagene Inc Tel 1-866-964-2589 Email: [info@antageneinc.com](mailto:info@antageneinc.com)