



Mouse Monoclonal Antibody **NQO1** conjugated to Sepharose Beads

CatalogNo: **ANT8039-S**

Size 200ul

Storage Store at 4 °C for frequent use

Description

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays.

NQO1 (ANT0071R) Rabbit mAb

Formulation: 50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Host Species

- Rabbit
- Human, Mouse, Rat,

Reactivity

- WB, IHC, IF, IP, ELISA

Applications

MW

- 31kD (Calculated)
 - IgG, Kappa
- 31kD (Observed)

Isotype

Recommended Dilution Ratios

IP

Basic Information

Clonality Monoclonal

Clone Number ANT0071R


Immunogen Information

Sequence	NAD(P)H dehydrogenase [quinone] 1
Specificity	Endogenous


Target Information

Gene name	>>Ubiquinone and other terpenoid-quinone biosynthesis;>>Metabolic pathways;>>Biosynthesis of cofactors;>>Pathways in cancer;>>Chemical carcinogenesis reactive oxygen species;>>Hepatocellular carcinoma;>>Fluid shear stress and atherosclerosis		
Protein Name	NQO1		
	Organism	Gene ID	UniProt ID
	Human	1728;	P15559;
	Mouse		Q64669;
Cellular Localization	Cytoplasm		
Tissue specificity	Colon,Liver,Pooled,		
Function	Catalytic activity:NAD(P)H + a quinone = NAD(P)(+) + a hydroquinone.,cofactor:FAD.,enzyme regulation:Inhibited by dicoumarol.,Function:The enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinons involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis.,induction:By dioxin.,mass spectrometry: PubMed:11735396,miscellaneous:Quinone reductase accepts electrons from both NADH and NADPH with equal efficiency.,polymorphism:The Ser-187 polymorphism may be linked to susceptibility to forms of cancers.,similarity:Belongs to the NAD(P)H dehydrogenase (quinone) family.,subunit:Homodimer.,		


Validation Data




Rat kidney was stained with anti-NQO1 (ANT0071R) rabbit antibody



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-NQO1 (ANT0071R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: HeLa Lane 2: NIH-3T3 Lane 3: Rat kidney Lane 4: HepG2
 Predicted band size: 31kDa Observed band size: 31kDa



Human kidney was stained with anti-NQO1 (ANT0071R) rabbit antibody



Mouse kidney was stained with anti-NQO1 (ANT0071R) rabbit antibody

For Research use only, not for diagnostics and clinical use
Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com