

www.antageneinc.com

Mouse Monoclonal Antibody CD31 conjugated to Sepharose Beads

CatalogNo: ANT8079-S

Size 200ul

Storage Store at 4 °C for frequent use

Description

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays.

CD31 (ANT0041R) Rabbit mAb

Formulation: 50% slurry in PBS pH 7.2 with 0.01mg NaN3a3 preservative.

Host Species Reactivity Applications

• Rabbit • Human, • WB,IHC,IF,IP,ELISA

MW Isotype

• 83kD (Calculated) • IgG,Kappa
130kD (Observed)

## **Recommended Dilution Ratios**

IP

### **Basic Information**

**Clonality** Monoclonal

Clone Number ANT0041R

# Immunogen Information Specificity

#### Endogenous

Gene name PECAM1

Protein Name Platelet endothelial cell adhesion molecule (PECAM-1) (EndoCAM) (GPIIA') (PECA1) (CD

antigen CD31)

Organism Gene ID UniProt ID

Human 5175; P16284;

Cellular Localization Membranous

Tissue specificity Expressed on platelets and leukocytes and is primarily concentrated at the borders between


endothelial cells (PubMed:18388311, PubMed:21464369). Expressed in human umbilical vein endothelial cells (HUVECs) (at protein level) (PubMed:19342684, PubMed:17580308). Expressed on neutrophils (at protein level) (PubMed:17580308). Isoform Long predominates in all tissues examined (PubMed:12433657). Isoform Delta12 is detected only in trachea (PubMed:12433657). Isoform Delta14-15 is only detected in lung (PubMed:12433657).

Isoform Delta14 is detected in all tissues examined with the strongest expression in heart (PubMed:12433657). Isoform Delta15 is expressed in brain, testis, ovary, cell surface of platelets, human umbilical vein endothelial cells (HUVECs), Jurkat T-cell leukemia, human erythroleukemia (HEL) and U-937 histiocytic lymphoma cell lines (at protein level) (PubMed:12433657,

PubMed:18388311).

**Function** 


Function:This protein is a cell adhesion molecule expressed on platelets and at endothelial cell intercellular junctions.,online information:CD31 entry,online information:PECAM-1,online information:The Singapore human mutation and polymorphism database,ANTM:Phosphorylated on Ser and Tyr residues after cellular activation.,similarity:Contains 6 Ig-like C2-type (immunoglobulin-like) domains.,tissue specificity:Long isoform predominates all tissues examined, isoform Delta12 was detected only in trachea and isoform Delta14-15 only in lung, isoform Delta14 was detected in all tissues examined with the strongest expression in heart.,




# **Validation**

### **Data**

Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-CD31 (ANT0041R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: THP-1 Predicted band size: 83kDa Observed band size: 130kDa



Human placenta was stained with Anti-CD31 (ANT0041R) rabbit antibody



Human tonsil was stained with Anti-CD31 (ANT0041R) rabbit antibody

For Research use only, not for diagnostics and clinical use
Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com