



Mouse Monoclonal Antibody **BRG1** conjugated to Sepharose Beads

CatalogNo: **ANT8093-M**

Size 200ul

Storage Store at 4 °C for frequent use

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

BRG1 (ANT0057R) Rabbit mAb

Formulation: Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg ANaN3.

Host Species

- Rabbit

Reactivity

- Human,Rat,

Applications

- WB,IHC,IF,IP,ELISA

MW

- 185kD (Calculated)
220kD (Observed)

Isotype

- IgG,Kappa

Recommended Dilution Ratios

IP

Basic Information

Clonality Monoclonal

Clone Number ANT0057R

Endogenous

Gene name SMARCA4
Protein Name Transcription activator BRG1

Organism	Gene ID	UniProt ID
Human	6597;	P51532;
Mouse	20586;	Q3TKT4;
Rat		Q8K1P7;


Cellular Nucleus

Localization


Tissue specificity Colocalizes with ZEB1 in E-cadherin-negative cells from established lines, and stroma of normal colon as well as in de-differentiated epithelial cells at the invasion front of colorectal carcinomas (at protein level).

Function Function:Transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. Also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene.,ANTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the SNF2/RAD54 helicase family.,similarity:Contains 1 bromo domain.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 HSA domain.,subunit:Interacts with NR3C1, PGR, SMARD1, TOPBP1 and ZMIM2/ZIMP7. Component of the BAF complex, which includes at least actin (ACTB), ARID1A, ARID1B/BAF250, SMARCA2, SMARCA4/BRG1, ACTL6A/BAF53, ACTL6B/BAF53B, SMARCE1/BAF57, SMARCC1/BAF155, SMARCC2/BAF170, SMARCB1/SNF5/INI1, and one or more of SMARCD1/BAF60A, SMARCD2/BAF60B, or SMARCD3/BAF60C. In muscle cells, the BAF complex also contains DPF3. Component of the BAF53 complex, at least composed of BAF53A, RUVBL1, SMARCA4/BRG1, and TRRAP, which preferentially acetylates histone H4 (and H2A) within nucleosomes. Component of the WINAC complex, at least composed of SMARCA2, SMARCA4, SMARCB1, SMARCC1, SMARCC2, SMARCD1, SMARCE1, ACTL6A, BAZ1B/WSTF, ARID1A, SUPT16H, CHAF1A and

Validation Data



Rat stomach was stained with anti-BRG1 (ANT0057R)



rabbit antibody

Various whole cell lysates were separated by 4-8% SDS-PAGE, and the membrane was blotted with anti-BRG1 (ANT0057R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: C6 Lane 2: HeLa Lane 3: HepG2
Predicted band size: 185kDa Observed band size: 220kDa