



LC3B (ANT0035R) Rabbit mAb

CatalogNo: ANT8147 **Recombinant** 

Formulation: PBS,50%glycerol,0.05%Proclin 300,0.05%BSA
Quantity : 100 ug/vial

Host Species

- Rabbit

MW

- 14kD,16kD (Calculated)
14kD,16kD (Observed)

Reactivity

- Human,Mouse,Rat,

Isotype

- IgG,Kappa

Applications

- WB,IHC,IF,IP,ELISA

Recommended Dilution Ratios

IHC 1:200-1:1000

WB 1:1000-1:5000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200,

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Basic Information

Clonality Monoclonal

Clone Number ANT0035R

Target Information

Immunogen Information Specificity

Endogenous

Gene name MAP1LC3B

Protein Name MAP1LC3B


Organism	Gene ID	UniProt ID
Human	81631 ;	Q9GZQ8 ;
Mouse		Q9CQV6 ;

Cellular Localization Cytoplasm

Tissue specificity Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver.


Function Caution:PubMed:12740394 has shown that the protein is cleaved at Lys-122 but PubMed:15355958 has shown that the cleavage site is at Gly-120 as in other mammalian orthologs.,Function:Probably involved in formation of autophagosomal vacuoles (autophagosomes).,ANTM:The precursor molecule is cleaved by APG4B/ATG4B to form LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form LC3-II.,similarity:Belongs to the MAP1 LC3 family.,subcellular location:LC3-II binds to the autophagic membranes.,subunit:3 different light chains, LC1, LC2 and LC3, can associate with MAP1A and MAP1B proteins.,tissue specificity:Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver.,

Validation Data









Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-LC3B (ANT0035R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1:Jurkat Lane 2:Rat spleen Lane 3: Mouse brain Lane 3: U-87 MG Predicted band size: 14,16kDa Observed band size: 14,16kDa

Mouse brain was stained with anti-LC3B (ANT0035R) rabbit antibody



Ret brain was stained with anti-LC3B (ANT0035R) rabbit antibody




Human brain was stained with anti-LC3B (ANT0035R) rabbit antibody

Jia, Wei-Qiang, et al. "Verbascoside inhibits progression of glioblastoma cells by promoting Let-7g-5p and down-regulating HMGA2 via Wnt/beta-catenin signalling blockade." *Journal of cellular and molecular medicine* 24.5 (2020): 2901-2916.

Immunofluorescence analysis of Hela cell. 1, Cleaved-PARP-1 (D214) Antibody(red) was diluted at 1:200(4° overnight). LC3B Antibody(green) was diluted at 1:200(4° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog:RS3611 was diluted at 1:1000(room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog:RS3208 was diluted at 1:1000(room temperature, 50min).

Immunofluorescence analysis of Mouse-lung tissue. 1, LC3B Antibody(red) was diluted at 1:200(4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B

Feng, Qin. "Quantitative proteomic analysis reveals that Arctigenin alleviates concanavalin A-induced hepatitis through suppressing immune system and regulating autophagy." *Frontiers in immunology* 9 (2018): 1881.



Western blot analysis of lysates from 1) HeLa, 2) 3T3, 3) Rat Brain Tissue cells, (Green) primary antibody was diluted at 1:1000, 4° over night, secondary antibody(cat:RS23920) was diluted at 1:10000, 37° 1hour. (Red) GAPDH Monoclonal Antibody(2B8) (cat:YM3029) antibody was diluted at 1:5000 as loading control, 4° over night, secondary antibody(cat:RS23710) was diluted at 1:10000, 37° 1hour.

For Research use only, not for diagnostics and clinical use
Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com