



Mouse Monoclonal Antibody **TSG101** conjugated to Sepharose Beads

CatalogNo: **ANT8150-M**

Size 200ul

Storage Store at 4 °C for frequent use

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

TSG101 (ANT0040R) Rabbit mAb

Formulation: Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg ANaN3.

Host Species

- Rabbit
- Human, Mouse, Rat,

Reactivity

- WB, IF, IP, ELISA

Applications

MW

- 44kD (Calculated)
- 44kD (Observed)
- IgG, Kappa

Isotype

Recommended Dilution Ratios

IP

Basic Information

Clonality Monoclonal

Clone Number ANT0040R

Immunogen Information

Specificity Endogenous

Target Information

Gene name TSG101
Protein Name Tumor susceptibility gene 101 protein


Organism	Gene ID	UniProt ID
Human	7251 ;	Q99816 ;
Mouse	22088 ;	Q61187 ;
Rat	292925 ;	Q6IRE4 ;

Cellular Localization Cytoplasm, Nuclear

Tissue specificity Heart, brain, placenta, lung, liver, skeletal, kidney and pancreas.

Function Alternative products:Additional isoforms seem to exist. Several shorter isoforms are detected in primary breast cancers and other tumors,Domain:The coiled coil domain may interact with stathmin.,Domain:The UEV domain binds ubiquitin and P-[ST]-A-P peptide motif independently.,Domain:The UEV domain is required for the interaction of the complex with ubiquitin. It also mediates the interaction with ANTAP/PSAP motifs of HIV-1 P6 protein and human spumaretrovirus Gag protein.,Function:Component of the ESCRT-I complex, a regulator of vesicular trafficking process. Binds to ubiquitinated cargo proteins and is required for the sorting of endocytic ubiquitinated cargos into multivesicular bodies (MVBs). Mediates the association between the ESCRT-0 and ESCRT-I complex. Required for completion of cytokinesis; the function requires CEP55. May be involved in cell growth and differentiation. Acts as a negative growth regulator. Involved in the budding of many viruses through an interaction with viral proteins that contain a late-budding motif P-[ST]-A-P. This interaction is essential for viral particle budding of numerous retroviruses.,PTM:Monoubiquitinated at multiple sites by LRSAM1. Ubiquitination inactivates it, possibly by regulating its shuttling between an active membrane-bound protein and an inactive soluble form.,similarity:Belongs to the ubiquitin-conjugating enzyme family. UEV subfamily.,similarity:Contains 1 SB (steadiness box) domain.,similarity:Contains 1 UEV (ubiquitin E2 variant) domain.,subcellular location:Mainly cytoplasmic. Membraneassociated when active and soluble when inactive. Depending on the stage of the cell cycle, detected in the nucleus. Colocalized with CEP55 in the midbody during cytokinesis.,subunit:Component of the ESCRT-I complex (endosomal sorting complex required for transport I) which consists of TSG101, VPS28, a VPS37 protein (VPS37A to -D) and a FAM125/MVB12 protein (FAM125A or -B) in a 1:1:1:1 stoichiometry. Interacts with VPS37A, VPS37B and VPS37C. Interacts with ubiquitin, stathmin, GMCL, DMAP1 and AATF (By similarity). Interacts with HGS; the interaction mediates the association with the ESCRT-0 complex. Interacts with GGA1 and GGA3. Interacts (via UEV domain) with PDCD6IP/AIP1. Interacts with VPS28, SNF8 and VPS36. Self-associates. Interacts with FAM125A/MVB12A; the association appears to be mediated by the TSG101-VPS37 binary subcomplex. Interacts with VPS37D. Interacts with LRSAM1. Interacts with CEP55; the interaction is required for cytokinesis but not for viral budding. Interacts with HIV-1 p6. Interacts with human spumavirus Gag. Interacts with HTLV-1 Gag. Interacts with Ebola virus VP40. Interacts with EIAV p9; the interaction has been shown in vitro.,tissue specificity:Heart, brain, placenta, lung, liver, skeletal, kidney and pancreas.,

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-TSG101 (ANT0040R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: Jurkat Lane 2: K562 Lane 3: NIH-3T3 Lane 4: PC-12

Predicted band size: 44kDa Observed band size: 44kDa

For Research use only, not for diagnostics and clinical use
Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com