



www.antageneinc.com

### Mouse Monoclonal Antibody SOX9 conjugated to Sepharose Beads

CatalogNo: ANT8170-M

Size 200ul

Storage Store at 4 °C for frequent use

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamidemodified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

## SOX9 (ANT0095R) Rabbit mAb

Formulation: Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg ANaN3.

Host Species <ul> <li>Rabbit</li> </ul>	• Human,Mouse,Rat,	<ul><li>Reactivity</li><li>WB,IHC,IF,ELISA</li></ul>	Applications
MW • 56kD (Calc 70kD (Obser	, , , , , , ,	Isotype	

## Recommended Dilution Ratios

#### IP

**Basic Information** 

Clonality Monoclonal

Clone Number ANT0095R


### Immunogen Information

Specificity Endogenous

# Target Information

Gene name Protein Name	SOX9 Transcription factor SOX-9							
	Orga	anism	Gen	e ID	UniProt ID			
		ł	luman	<u>6662</u> ;	<u>P48436</u> ;			
	Cellular Localization	Nucleus						
	Tissue specificity Eye, PNS, Testis,							
	Function	Disease:Defects in SOX9 are the cause of campomelic dysplasia (CMD1) [MIM:114290]. CMD1 is a rare, often lethal, dominantly inherited, congenital osteochondrodysplasia, associated with male-to-female autosomal sex reversal in two-thirds of the affected karyotypic males. A disease of the newborn characterized by congenital bowing and angulation of long bones, unusually small scapulae, deformed pelvis and spine and a missing pair of ribs. Craniofacial defects such as cleft palate, micrognatia, flat face and hypertelorism are common. Various defects of the ear are often evident, affecting the cochlea, malleus incus, stapes and tympanum. Most patients die soon after birth due to respiratory distress which has been attributed to hypoplasia of the tracheobronchial cartilage and small thoracic cage.,Function:Plays an important role in the normal skeletal development. May regulate the expression of other genes involved in chondrogenesis by acting as a transcription factor for these genes.,similarity:Contains 1 HMG box DNA-binding domain.,						

## Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-SOX9 (ANT0095R) antibody. The HRPconjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HepG2 Lane 2: U-14 Lane 3: SW480 Predicted band size: 56kDa Observed band size: 70kDa

For Research use only, not for diagnostics and clinical use Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com