



Mouse Monoclonal Antibody **PPAR α** conjugated to Sepharose Beads

CatalogNo: **ANT8234-M**

Size 200ul

Storage Store at 4 °C for frequent use

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

PPAR α (ANT0084R) Rabbit mAb

Formulation: Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg ANaN3.

Host Species

- Rabbit
- Human, Mouse, Rat,

Reactivity

- WB, IHC, IF, IP, ELISA

Applications

MW

- 52kD (Calculated)
- 52kD (Observed)
- IgG, Kappa

Isotype

Recommended Dilution Ratios

IP

Basic Information

Clonality Monoclonal

Clone Number ANT0084R

Immunogen Information

Specificity Endogenous

Gene name PPARA

Protein Name Peroxisome proliferator-activated receptor alpha


Organism	Gene ID	UniProt ID
Human	5465;	Q07869;
Mouse	19013;	P23204;
Rat	25747;	P37230;

Cellular Localization Nucleus


Tissue specificity Skeletal muscle, liver, heart and kidney. Expressed in monocytes (PubMed:28167758).

Function Function:Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the receptor binds to a promoter element in the gene for acyl-CoA oxidase and activates its transcription. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids.,online information:Peroxisome proliferator-activated receptor entry,similarity:Belongs to the nuclear hormone receptor family. NR1 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding domain.,subunit:Heterodimer with the retinoid X receptor. Interacts with NCOA3 and NCOA6 coactivators, leading to a strong increase of transcription of target genes. Also interacts with PPARBP coactivator in vitro. Interacts with AKAP13.,tissue specificity:Skeletal muscle, liver, heart and kidney.,


Validation Data




Rat liver was stained with anti-PPARα (ANT0084R) rabbit antibody




Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-PPARα (ANT0084R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: HepG2 Lane 2: MCF7 Lane 3: NIH-3T3 Lane 4: L6 Predicted band size: 52kDa Observed band size: 52kDa




Human hepatocellular carcinoma was stained with anti-PPAR α (ANT0084R) rabbit antibody



Human liver was stained with anti-PPAR α (ANT0084R) rabbit antibody



Mouse liver was stained with anti-PPAR α (ANT0084R) rabbit antibody



Immunofluorescence analysis of HeLa cell. 1, PPAR- α Antibody (red) was diluted at 1:200 (4° overnight). Galectin-3 Monoclonal Antibody (6G2) (green) was diluted at 1:200 (4° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog: RS3611 was diluted at 1:1000 (room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog: RS3208 was diluted at 1:1000 (room temperature, 50min).

For Research use only, not for diagnostics and clinical use
Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com