



www.antageneinc.com

Mouse Monoclonal Antibody CDK4 conjugated to Sepharose Beads

CatalogNo: ANT8244-M

Size 200ul

Storage Store at 4 °C for frequent use

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

CDK4 (ANT0099R) Rabbit mAb

Formulation: Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg ANaN3.

Host Species Reactivity Applications

Rabbit
 Human, Mouse, Rat,
 WB, IHC, IF, IP, ELISA

MW Isotype

• 34kD (Calculated) • IgG,Kappa

34kD (Observed)

Recommended Dilution Ratios

ID

Basic Information

Clonality Monoclonal

Clone Number ANT0099R

Immunogen Information

Specificity Endogenous

Target Information

Gene name

CDK4

Protein Name

Cyclin-dependent kinase 4

Organism	Gene ID	UniProt ID
Human	<u>1019</u> ;	<u>P11802</u> ;
Mouse	<u>12567</u> ;	<u>P30285</u> ;
Rat	94201;	<u>P35426</u> ;


Cellular Localization Cytoplasm, Nucleus

Tissue specificity Brain, Muscle,

Function


Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,Disease:CDK4 mutations are involved in tumor formation.,Disease:Defects in CDK4 are the cause of cutaneous malignant melanoma 3 (CMM3) [MIM:609048, 155600]. Malignant melanoma is a malignant neoplasm of melanocytes, arising de novo or from a preexisting benign nevus, which occurs most often in the skin but also may involve other sites.,enzyme regulation:Phosphorylation at Thr-172 is necessary for enzymatic activity.,Function:Probably involved in the control of the cell cycle.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Forms a stable complex with D-type G1 cyclins. Interacts with SEI1 and ZNF655/VIK.,

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-CDK4 (ANT0099R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: Hela Lane 2: C6 Lane 3: 3T3-L1 Predicted band size: 34kDa

Observed band size: 34kDa



Immunofluorescence analysis of Hela cell. 1,Cdk4 Antibody(green) was diluted at 1:200(4° overnight). 2, Goat Anti Rabbit Alexa Fluor 488 Catalog:RS3211 was diluted at 1:1000(room temperature, 50min). 3 DAPI(blue) 10min.

For Research use only, not for diagnostics and clinical use
Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com