



www.antageneinc.com

Mouse Monoclonal Antibody CD14 conjugated to Sepharose Beads

CatalogNo: ANT8259-M

Size 200ul

Storage Store at 4 °C for frequent use

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

CD14 (ANT0017R) Rabbit mAb

Formulation: Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg ANaN3.

Host Species Reactivity Applications

Rabbit
 Human, Mouse,
 WB, IHC, IF, IP, ELISA

MW Isotype

• 39kD (Calculated) • IgG,Kappa

52kD (Observed)

Recommended Dilution Ratios

IP

Basic Information

Clonality Monoclonal

Clone Number ANT0017R

Immunogen Information

Specificity Endogenous

Gene name CD14
Protein Name CD14

| Organism | Gene ID | UniProt ID |
|----------|----------------|-----------------|
| Human | <u>929</u> ; | <u>P10810</u> ; |
| Mouse | <u>12475</u> ; | <u>P10810</u> ; |
| Rat | <u>60350</u> ; | <u>Q63691</u> ; |

Cellular

Membrane

Localization

Tissue specificity Detected on macrophages (at protein level) (PubMed:1698311). Expressed strongly on the surface of monocytes and weakly on the surface of granulocytes; also expressed by most tissue macrophages.

Function Function: Cooperates with MD-2 and TLR4 to mediate the innate immune response to bacterial

lipopolysaccharide (LPS). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Up-regulates cell surface molecules, including adhesion

molecules., online information: CD14 entry, similarity: Contains 11 LRR (leucine-rich)


repeats., subunit: Belongs to the lipopolysaccharide (LPS) receptor, a multi-protein complex containing

at least CD14, MD-2 and TLR4., tissue


specificity:Expressed strongly on the surface of monocytes and weakly on the surface of granulocytes;

also expressed by most tissue macrophages.,


Validation Data



Mouse spleen was stained with anti-CD14 (ANT0017R) rabbit antibody



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-CD14 (ANT0017R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: RAW264.7 Predicted band size: 39kDa Observed band size: 52kDa



Human tonsil was stained with anti-CD14 (ANT0017R) rabbit antibody

For Research use only, not for diagnostics and clinical use
Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com