



## MIF (ANT0018R) Rabbit mAb

CatalogNo: ANT8260 **Recombinant** 

Formulation: PBS,50%glycerol,0.05%Proclin 300,0.05%BSA  
Quantity : 100 ug/vial

### Host Species

- Rabbit
- Human,Mouse,Rat,

### Reactivity

- WB,IHC,IF,IP,ELISA

### Applications

### MW

- 13kD (Calculated)
- 13kD (Observed)

### Isotype

- IgG,Kappa

## Recommended Dilution Ratios

IHC 1:3000-1:10000

WB 1:1000-1:5000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

## Storage

**Storage\*** -15°C to -25°C/1 year(Do not lower than -25°C)

## Basic Information

**Clonality** Monoclonal

**Clone Number** ANT0018R

## Target Information

# Immunogen Information Specificity

Endogenous

**Gene name** MIF  
**Protein Name** Macrophage migration inhibitory factor

Organism	Gene ID	UniProt ID
Human	<a href="#">4282;</a>	<a href="#">P14174;</a>
Mouse	<a href="#">17319;</a>	<a href="#">P34884;</a>
Rat	<a href="#">81683;</a>	<a href="#">P30904;</a>


**Cellular Localization** Cytoplasm

**Tissue specificity** Brain,Lens,Liver,Lung,Platelet,Skin

**Function** Catalytic activity:Keto-phenylpyruvate = enol-phenylpyruvate.,Disease:Genetic variations in MIF are associated with susceptibility to systemic juvenile rheumatoid arthritis [MIM:604302]. Systemic juvenile rheumatoid arthritis is juvenile chronic arthritis associated with severe, debilitating, extraarticular features, and occasionally fatal complications. Despite medical treatment, many children still experience early joint destruction, necessitating surgical replacement.,Function:The expression of MIF at sites of inflammation suggest a role for the mediator in regulating the function of macrophage in host defense. Also acts as a phenylpyruvate tautomerase.,similarity:Belongs to the MIF family.,subunit:Homotrimer. Interacts with COPS5 and BNIPL.,

---

## Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-MIF (ANT0018R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: C6 Lane 2: 3T3-L1 Lane 3: Jurkat Lane 4: Y79 Predicted band size: 13kDa Observed band size: 13kDa

Human thymus was stained with anti-MIF (ANT0018R) rabbit antibody

Immunofluorescence analysis of A549. 1,primary Antibody was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 488 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, Picture B: DAPI(blue) 10min.

Immunofluorescence analysis of HeLa cell. 1,MIF Antibody(green) was diluted at 1:200(4° overnight). 2, Goat Anti Rabbit Alexa Fluor 488 Catalog:RS3211 was diluted at 1:1000(room temperature, 50min). 3 DAPI(blue) 10min.

For Research use only, not for diagnostics and clinical use  
Contact Antagene Inc Tel 1-866-964-2589 Email: [info@antageneinc.com](mailto:info@antageneinc.com)