

www.antageneinc.com

CD20 (ANT0019R) Rabbit mAb

CatalogNo: ANT8261 Recombinant R

Formulation: PBS,50%glycerol,0.05%Proclin 300,0.05%BSA

Quantity: 100 ug/vial

Host Species Reactivity Applications

Rabbit
 Human,
 WB,IHC,IF,IP,ELISA

MW Isotype

• 33kD (Calculated) • IgG,Kappa

36kD (Observed)

Recommended Dilution Ratios

IHC 1:500-1:5000 WB 1:1000-1:5000 IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Basic Information

Clonality Monoclonal

Clone Number ANT0019R

Target Information

Immunogen Information Specificity

Endogenous

Gene name MS4A1 CD20

Protein Name CD20

Organism	Gene ID	UniProt ID
Human	<u>931</u> ;	<u>P11836</u> ;
Mouse	<u>12482</u> ;	<u>P19437</u> ;
Rat	<u>309217</u> ;	<u>D4A4Y3</u> ;


Cellular Localization Membrane

Tissue specificity Expressed on B-cells.


Function

Caution:Epitope 1, mapped in PubMed:16785532, is predicted to be buried in the membrane. Its accessibility to the extracellular space, and thus to antibody recognition, is not explained., Function:This protein may be involved in the regulation of B-cell activation and proliferation., online information:CD20 entry, pharmaceutical:Monoclonal antibodies (mAb) against CD20 are used to treat B-cell non-Hodgkin's lymphoma (NHL). These antibodies include Rituximab (Mabthera), Britumomab (Zevalin) and Tositumomab (Bexxar). CD20 engaged by mAb can generate transmembrane signals capable of directly controlling cell growth and triggering cell death in certain tumors. Alternatively, mAb can mediate complement-dependent cytotoxicity., ANTM:Phosphorylated. Might be functionally regulated by protein kinase(s)., similarity:Belongs to the MS4A family., tissue specificity:Expressed on B-cells.,


Validation Data



Human tonsil was stained with anti-CD20 (ANT0019R) rabbit antibody



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-CD20 (ANT0019R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: Raji Lane 2: Daudi Predicted band size: 33kDa Observed band size: 36kDa



Human spleen was stained with anti-CD20 (ANT0019R) rabbit antibody

For Research use only, not for diagnostics and clinical use
Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com