



Mouse Monoclonal Antibody **RUNX2** conjugated to Sepharose Beads

CatalogNo: **ANT8347-S**

Size 200ul

Storage Store at 4 °C for frequent use

Description

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays.

RUNX2 (ANT0024R) Rabbit mAb

Formulation: 50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Host Species

- Rabbit
- Human, Mouse, Rat,

Reactivity

- WB, IHC, IF, IP, ELISA

Applications

MW

- 57kD (Calculated)
 - IgG, Kappa
- 57kD (Observed)

Isotype

Recommended Dilution Ratios

IP

Basic Information

Clonality

Monoclonal

Clone Number ANT0024R

Immunogen Information

Specificity Endogenous

Gene name RUNX2

Protein Name Runt-related transcription factor 2


Organism	Gene ID	UniProt ID
Human	860 ;	Q13950 ;
Mouse	12393 ;	Q08775 ;
Rat		Q9Z2J9 ;

Cellular Localization Nucleus

Tissue specificity Specifically expressed in osteoblasts.

Function Disease:Defects in RUNX2 are the cause of cleidocranial dysplasia (CCD) [MIM:119600]. CCD is an autosomal dominant skeletal disorder with high penetrance and variable expressivity. It is due to defective endochondral and intramembranous bone formation. Typical features include hypoplasia/aplasia of clavicles, patent fontanelles, wormian bones (additional cranial plates caused by abnormal ossification of the calvaria), supernumerary teeth, short stature, and other skeletal changes. In some cases defects in RUNX2 are exclusively associated with dental anomalies.,Domain:A proline/serine/threonine rich region at the C-terminus is necessary for transcriptional activation of target genes and contains the phosphorylation sites.,Function:Transcription factor involved in osteoblastic differentiation and skeletal morphogenesis. Essential for the maturation of osteoblasts and both intramembranous and endochondral ossification. CBF binds to the core site, 5'PYGPYGGT-3', of a number of enhancers and promoters, including murine leukemia virus, polyomavirus enhancer, T-cell receptor enhancers, osteocalcin, osteopontin, bone sialoprotein, alpha 1(I) collagen, LCK, IL-3 and GM-CSF promoters (By similarity). Inhibits MYST4-dependent transcriptional activation.,ANTM:Phosphorylated; probably by MAP kinases (MAPK) (By similarity). Isoform 3 is phosphorylated on Ser-340.,similarity:Contains 1 Runt domain.,subunit:Heterodimer of an alpha and a beta subunit. Interacts with HIVEP3 (By similarity). The alpha subunit binds DNA as a monomer and through the Runt domain. DNAbinding is increased by heterodimerization. Interacts with G22P1 (Ku70) and XRCC5 (Ku80). Interacts with MYST3 and MYST4.,tissue specificity:Specifically expressed in osteoblasts.,

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-RUNX2 (ANT0024R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: MDA-MB-231 Lane 2: Saos-2 Lane 3: NIH-3T3 Lane 4: Rat womb Predicted band size: 57kDa Observed band size: 57kDa Immunofluorescence analysis of HeLa cell. 1, RUNX2 Antibody(red) was diluted at 1:200(4° overnight). NSE Monoclonal Antibody(13E2)(green) was diluted at 1:200(4° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog:RS3611 was diluted at 1:1000(room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog:RS3208 was diluted at 1:1000(room temperature, 50min).

For Research use only, not for diagnostics and clinical use
Contact Antagene Inc Tel 1-866-964-2589 Email: info@antageneinc.com