



## Mouse Monoclonal Antibody **CREBBP** conjugated to Sepharose Beads

CatalogNo: **ANT8348-M**

Size 200ul

Storage Store at 4 °C for frequent use

### Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

### **CREBBP (ANT0026R) Rabbit mAb**

Formulation: Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg ANaN3.

#### Host Species

- Rabbit

#### MW

- 300kD (Calculated)
- 300kD (Observed)

#### Reactivity

- Human,Mouse,Rat,

#### Isotype

- IgG,Kappa

#### Applications

- WB,IHC,IF,IP,ELISA

## **Recommended Dilution Ratios**

### **IP**

## **Basic Information**

#### Clonality

Monoclonal

Clone Number      ANT0026R

**Immunogen Information**

**Specificity**      Endogenous

**Gene name**      CREBBP CBP

**Protein Name**      CREB-binding protein (EC 2.3.1.48)


Organism	Gene ID	UniProt ID
Human	<a href="#">1387</a> ;	<a href="#">Q92793</a> ;
Mouse		<a href="#">P45481</a> ;

**Cellular  
Localization**      Nucleus

**Function**      Acetylates histones, giving a specific tag for transcriptional activation . Also acetylates nonhistone proteins, like DDX21, FBL, IRF2, MAFG, NCOA3, POLR1E/PAF53 and FOXO1 . Binds specifically to phosphorylated CREB and enhances its transcriptional activity toward cAMPResponsive genes. Acts as a coactivator of ALX1. Acts as a circadian transcriptional coactivator which enhances the activity of the circadian transcriptional activators: NPAS2ARNTL/BMAL1 and CLOCK-ARNTL/BMAL1 heterodimers . Acetylates PCNA; acetylation promotes removal of chromatin-bound PCNA and its degradation during nucleotide excision repair (NER) . Acetylates POLR1E/PAF53, leading to decreased association of RNA polymerase I with the rDNA promoter region and coding region . Acetylates DDX21, thereby inhibiting DDX21 helicase activity . Acetylates FBL, preventing methylation of 'Gln-105' of histone H2A (H2AQ104me) . Functions as a transcriptional coactivator for SMAD4 in the TGF-beta signaling pathway .


---

## Validation Data



Various whole cell lysates were separated by 4-8% SDS-PAGE, and the membrane was blotted with anti-CREBBP (ANT0026R) antibody. The HRPconjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: HeLa Predicted band size: 300kDa Observed band size: 300kDa

Human pancreas was stained with anti-CREBBP (ANT0026R) rabbit antibody



Mouse colon was stained with anti-CREBBP (ANT0026R) rabbit antibody

---

For Research use only, not for diagnostics and clinical use  
Contact Antagene Inc Tel 1-866-964-2589 Email: [info@antageneinc.com](mailto:info@antageneinc.com)