



**Category:** Polyclonal Antibodies  
**Cat. #** 1107NF-103M

**Product Name:** Polyclonal Antibody Goat Anti-Human IgM (Mu chains) – Prediluted

**Description:**  
Polyclonal Antibody Goat Anti-Human IgM (Mu chains)

**Immunogen:**  
Purified human IgM Mu chains.

**Application:**  
This antibody is prediluted for Immunohistochemistry use.

**Species Reactivity:**  
Human. Others not tested.

**Recommended Positive Control:**  
Tonsil, Spleen

**Presentation:**  
Polyclonal antibody prediluted in Primary Antibody Diluting Buffer pH 7.6 and quality controlled using formalin-fixed paraffin-embedded human lymphoid tissue sections.

**Aliquoting Instructions:**  
Not applicable

**Staining Procedure:**  
The prediluted antibody can be used on frozen cryostat sections as well as formalin-fixed paraffin-embedded tissue sections. For paraffinembedded tissue sections, we recommend an incubation time and temperature of 30 minutes at 37°C for this antibody, when used in conjunction with an immunoperoxidase staining kit.  
\*Prolonged fixation in buffered formalin can destroy the epitope.

**Specificity:**  
The antibody reacts specifically with IgM immunoglobulins.

**Storage:**  
Refrigerate at 4°C. Do not freeze.

**Size:** 5 ml  
**Host:** Goat  
**Form:** Prediluted  
**Concentration:** .007-.010 mg/ml  
**Units On Hand:** YES

**For Research Use Only**

**Contact:** Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: [Info@antageneinc.com](mailto:Info@antageneinc.com)