



Product Information Sheet

Monoclonal Anti- α -Fetoprotein, AFP conjugated to Magnetic Beads

Catalogue No. MA1001-M

Lot No. 08A12

Clone: AP-1

Ig type: mouse IgG2a

Size: 200 μ l

Specificity

Human.

No cross reactivity with other proteins.

Recommended application

Immunoprecipitation(IP)



Immunogen

Human α -fetoprotein

Purification

Purified by the goat anti-mouse IgG affinity chromatography.

Formulation

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN₃.

Storage

Store at 4°C for frequent use.

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

BACKGROUND

Alpha-fetoprotein (AFP) is a major plasma protein in the fetus, where it is produced by the yolk sac and liver. Direct confirmation of the assignment of the AFP gene to chromosome 4 by in situ hybridization was provided by Harper and Dugaiczky (1983), who placed the gene in the q11-q22 region, the same region as the albumin gene. Structure and evolution of human alpha-fetoprotein deduced from partial sequence of cloned cDNA. As the major fetal serum protein, Alpha-fetoprotein is not essential for embryonic development but is required for female fertility.

REFERENCE

1. D'Eustachio, P.; Ingram, R. S.; Tilghman, S. M.; Ruddle, F. H. : Murine alpha-fetoprotein and albumin: two evolutionarily linked proteins encoded on the same mouse chromosome. *Somat. Cell Genet.* 7: 289-294, 1981.
2. Beattie, W. G.; Dugaiczky, A. : Structure and evolution of human alpha-fetoprotein deduced from partial sequence of cloned cDNA. *Gene* 20: 415-422, 1982.
3. Gabant, P.; Forrester, L.; Nichols, J.; Van Reeth, T.; De Mees, C.; Pajack, B.; Watt, A.; Smits, J.; Alexandre, H.; Szpirer, C.; Szpirer, J. : Alpha-fetoprotein, the major fetal serum protein, is not essential for embryonic development but is required for female fertility. *Proc. Nat. Acad. Sci.* 99: 12865-12870, 2002.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com



For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com