

## **Product Information Sheet**



Monoclonal Anti-α-Fetoprotein, AFP (Sepharose Bead Conjugate)

Catalogue No. MA1001-S	
	Immunogen
Lot No. 08A12	Human α-fetoprotein
Clone: AP-1	Purification
	Purified by the goat anti-mouse IgG affinity chromatography.
<b>Ig type:</b> mouse IgG2a	
	Formulation
<b>Size:</b> 200µl	50% slurry in PBS pH 7.2 with 0.01mg NaN $_3a_3$ preservative.
Specificity	Storage
Human.	Store at 4°C for frequent use.
No cross reactivity with other	
proteins.	Description:
	This Antagene antibody is immobilized via covalent binding of primary amino groups to
Recommended application	N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for
Immunoprecipitation(IP)	immunoprecipitation assays

## BACKGROUND

Alpha-fetoprotein (AFP) is a major plasma protein in the fetus, where it is produced by the yolk sac and liver. Direct confirmation of the assignment of the AFP gene to chromosome 4 by in situ hybridization was provided by Harper and Dugaiczyk (1983), who placed the gene in the q11-q22 region, the same region as the albumin gene. Structure and evolution of human alpha-fetoprotein deduced from partial sequence of cloned cDNA.As the major fetal serum protein, Alpha-fetoprotein is not essential for embryonic development but is required for female fertility.

## REFERENCE

1. D'Eustachio, P.; Ingram, R. S.; Tilghman, S. M.; Ruddle, F. H. : Murine alpha-fetoprotein and albumin: two evolutionarily linked proteins encoded on the same mouse chromosome. *Somat. Cell Genet.* 7: 289-294, 1981.

2. Beattie, W. G.; Dugaiczyk, A. : Structure and evolution of human alpha-fetoprotein deduced from partial sequence of cloned cDNA. *Gene* 20: 415-422, 1982.

3. Gabant, P.; Forrester, L.; Nichols, J.; Van Reeth, T.; De Mees, C.; Pajack, B.; Watt, A.; Smitz, J.; Alexandre, H.; Szpirer, C.; Szpirer, J. : Alpha-fetoprotein, the major fetal serum protein, is not essential for embryonic development but is required for female fertility. *Proc. Nat. Acad. Sci.* 99: 12865-12870, 2002.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com