



Product Information Sheet

Monoclonal Anti- Catenin γ (Plakoglobin) (Sepharose Bead Conjugate)

Catalogue No. MA1012-S

Immunogen

Recombinant chicken plakoglobin.

Lot No. 08A12

Purification

Clone: CN-3

Purified by the goat anti-mouse IgG affinity chromatography.

Ig type: mouse IgG1

Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Size: 200 μ l

Storage

Specificity

Human, rat, chicken.

No cross reactivity with other proteins.

Store at 4°C for frequent use.

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays.

Recommended application

Immunoprecipitation(IP)

BACKGROUND

Junction Plakoglobin (JUP) ,also knows as catenin gamma, is a major cytoplasmic protein that occurs in a soluble and a membrane-associated form and is the only known constituent common to the submembranous plaques of both kinds of adhering junctions, the desmosomes and the intermediate junctions. It is a component of the cadherin-catenin complex, which is predominantly localized where actin filaments anchor in adherens junctions of epithelial cells. The human plakoglobin gene localizes on chromosome 17q21. Gamma-catenin is regulated by the APC tumor suppressor and its oncogenic activity is distinct from that of beta-catenin.

REFERENCE

- 1 Aberle, H.; Bierkamp, C.; Torchard, D.; Serova, O.; Wagner, T.; Natt, E.; Wirsching, J.; Heidkamper, C.; Montagna, M.; Lynch, H. T.; Lenoir, G. M.; Scherer, G.; Feunteun, J.; Kemler, R. : The human plakoglobin gene localizes on chromosome 17q21 and is subjected to loss of heterozygosity in breast and ovarian cancers. *Proc. Nat. Acad. Sci.* 92: 6384-6388, 1995.
- 2 Kolligs, F. T.; Kolligs, B.; Hajra, K. M.; Hu, G.; Tani, M.; Cho, K. R.; Fearon, E. R. : Gamma-catenin is regulated by the APC tumor suppressor and its oncogenic activity is distinct from that of beta-catenin. *Genes Dev.* 14: 1319-1331, 2000.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com