



Product Information Sheet

Monoclonal Anti-CDC25C, conjugated to Magnetic Beads

Catalogue No. MA1018-M

Immunogen

Recombinant human Cdc25c.

Lot No. 08A12

Purification

Clone: IMD-25

Purified by the goat anti-mouse IgG affinity chromatography.

Ig type: mouse IgG1

Formulation

Size: 200µl

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN₃.

Specificity

Human.

No cross reactivity with other proteins.

Storage

Store at 4°C for frequent use.

Recommended application

Immunoprecipitation(IP)

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

BACKGROUND

Cell division cycle(CDC25) gene product is a protein-tyrosine phosphatase, activates a partially purified p34(cdc2)/cyclin B complex. The cdc25 protein also shares homology with a protein phosphatase with activity against both tyrosine and serine (and thus probably threonine) phosphate residues. CDC25C is expressed predominantly in G2 phase in healer cells. The human gene encodes a protein with a predicted molecular mass of 53,000 daltons whose C-terminal domain shares about 37% sequence identity with the fission yeast cdc25 mitotic inducer. CDC25C gene is mapped to 5q31.

REFERENCE

1. Sartor, H.; Ehlert, F.; Grzeschik, K.-H.; Muller, R.; Adolph, S. : Assignment of two human cell cycle genes, CDC25C and CCNB1, to 5q31 and 5q12, respectively. *Genomics* 13: 911-912, 1992.
2. Taviaux, S. A.; Demaille, J. G. : Localization of human cell cycle regulatory genes CDC25C to 5q31 and WEE1 to 11p15.3-11p.15.1 by fluorescence in situ hybridization. *Genomics* 15: 194-196, 1993.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com