

Product Information Sheet



Monoclonal Anti-c-Myc (Sepharose Bead Conjugate)

Catalogue No. MA1028-S	proteins.
Lot No. 08A12	Recommended application Immunoprecipitation(IP)
Clone: IMD-3	
Ig type: mouse IgG2a	
Size : 200µl	
Specificity	

Human.

sepharose beads. It is useful for immunoprecipitation assays.

No cross reactivity with other

BACKGROUND

C-Myc is an oncogene that functions both in the stimulation of cell proliferation and in apoptosis. c-Myc elicits its oncogenic activity by causing immortalization, and to a lesser extent the transformation of cells, in addition to several other mechanisms. The c-MYC proto-oncogene encodes a transcription factor that is critical for cell growth and proliferation. It is one of the genes frequently altered in cancer cells in which it exhibits constitutive activity. Downregulation of c-Myc is critical for 2-Methoxyestradiol (2ME2)-induced oxidative stress and apoptosis in AML cells. And its up-regulation is important for promoting lymphocyte cell division, and demonstrating that GFP-c-Myc expression is a marker of proliferating lymphocytes in vivo.

REFERENCE

- 1. Liao D, Thakur A, Wu J, Biliran H, Sarkar FH. Perspectives on c-Myc, Cyclin D1, and Their Interaction in Cancer Formation, Progression, and Response to Chemotherapy.Crit Rev Oncog. 2007;13(2):93-158.
- 2. Britton S, Salles B, Calsou P. c-MYC protein is degraded in response to UV irradiation. Cell Cycle. 2007 Oct 2;7(1)
- 3. Chow JM, Liu CR, Lin CP, Lee CN, Cheng YC, Lin S, Liu HE. Downregulation of c-Myc determines sensitivity to 2-methoxyestradiol-induced apoptosis in human acute myeloid leukemia.Exp Hematol. 2008 Feb;36(2):140-8.
- 4. Huang CY, Bredemeyer AL, Walker LM, Bassing CH, Sleckman BP. Dynamic regulation of c-Myc proto-oncogene expression during lymphocyte development revealed by a GFP-c-Myc knock-in mouse.Eur J Immunol. 2008 Jan 14.