



Product Information Sheet

Monoclonal Anti-Growth Associated Protein-43, GAP43 (Sepharose Bead Conjugate)

Catalogue No. MA1042-S Immunogen

GAP-43 from neonatal rat forebrain membranes.

Lot No. 08A12

Purification

Clone: GAP-8A12 Purified by the goat anti-mouse IgG affinity chromatography.

Ig type: mouse IgG2a Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃a₃ preservative.

Size: 200µl

Storage

Specificity Store at 4°C for frequent use.

Human, mouse, rat, chicken, snake

No cross reactivity with other

proteins.

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated

sepharose beads. It is useful for immunoprecipitation assays.

Recommended application

Immunoprecipitation(IP)

BACKGROUND

GAP43 is expressed by developing and regenerating neurons, and to a lesser extent, reactive glial cells. It is used widely to specifically label injured neurons and to score neuronal regeneration. GAP43 is also a neuronal growth cone protein thought to be involved in pathfinding. GAP43 is considered to be a crucial component of an effective regenerative response in the nervous system.

REFERENCE

1.Kosik, K. S.; Orecchio, L. D.; Bruns, G. A. P.; Benowitz, L. I.; MacDonald, G. P.; Cox, D. R.; Neve, R. L.: Human GAP-43: its deduced amino acid sequence and chromosomal localization in mouse and human. *Neuron* 1: 127-132, 1988.

2. Strittmatter, S. M.; Fankhauser, C.; Huang, P. L.; Mashimo, H.; Fishman, M. C.:

Neuronal pathfinding is abnormal in mice lacking the neuronal growth cone protein GAP-43. *Cell* 80: 445-452, 1995.