



Product Information Sheet

Monoclonal Anti-Glial Fibrillary Acidic Protein, *GFAP* - conjugated to Magnetic Beads

Catalogue No. MA1045-M

Immunogen

GFAP from pig spinal cord.

Lot No. 08A12

Purification

Clone: GA-8

Purified by the goat anti-mouse IgG affinity chromatography.

Ig type: mouse IgG1

Formulation

Size: 200µl

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN₃.

Specificity

Human, mouse, rat.

No cross reactivity with other proteins.

Storage

Store at 4°C for frequent use.

Recommended application

Immunoprecipitation(IP)

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

BACKGROUND

Glial fibrillary acidic protein(GFAP) is an intermediate filament protein of 52Kda. GFAP gene is mapped to human 17q21. GFAP is a useful marker of astroglia in the brain. Mutations in GFAP, encoding glial fibrillary acidic protein, are associated with Alexander disease

REFERENCE

- 1 Brenner, M.; Johnson, A. B.; Boespflug-Tanguy, O.; Rodriguez, D.; Goldman, J. E.; Messing, A. : Mutations in GFAP, encoding glial fibrillary acidic protein, are associated with Alexander disease. Nature Genet. 27: 117-120, 2001.
- 2 Rodriguez, D.; Gauthier, F.; Bertini, E.; Bugiani, M.; Brenner, M.; N'guyen, S.; Goizet, C.; Gelot, A.; Surtees, R.; Pedespan, J.-M.; Hernandorena, X.; Troncoso, M.; Uziel, G.; Messing, A.; Ponsot, G.; Pham-Dinh, D.; Dautigny, A.; Boespflug-Tanguy, O. : Infantile Alexander disease: spectrum of GFAP mutations and genotype-phenotype correlation. Am. J. Hum. Genet. 69: 1134-1140, 2001. Note: Erratum: Am. J. Hum. Genet. 69: 1413 only, 2001.

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC AND CLINICAL USE.