



Product Information Sheet

Monoclonal Anti-N-Cadherin (Sephacrose Bead Conjugate)

Catalogue No. MA1067-S

Lot No. 08A12

Clone: NC-17

Ig type: mouse IgG1

Size: 200µl

Specificity

Human, mouse, rat, rabbit, chicken.

No cross reactivity with other proteins.

Recommended application

Immunoprecipitation (IP)

Immunogen

Affinity purified chicken heart A-CAM.

Purification

Purified by the goat anti-mouse IgG affinity chromatography.

Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Storage

Store at 4°C for frequent use.

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays

BACKGROUND

N-cadherin (NCAD) is a member of the cadherin cell-cell adhesion receptor family that includes P-, E-, and R-cadherin and liver cell adhesion molecule (L-CAM). N-Cadherin, also known as Cadherin-2, encodes a 907-amino acid protein that includes a 159-amino acid signal sequence. Human and mouse nucleotide sequences are 96% identical. Mouse Ncad gene consists of 16 exons dispersed over more than 200 kb of genomic DNA. Human N-cadherin gene contains 16 exons and its sequence is highly similar to both the mouse NCAD gene (including the large first and second introns) and other cadherin genes. N-cadherin is mapped to 18q11.2. Cadherin regulates dendritic spine morphogenesis.

REFERENCE

- 1 Miyatani, S.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Takeichi, M. : Genomic structure and chromosomal mapping of the mouse N-cadherin gene. *Proc. Nat. Acad. Sci.* 89: 8443-8447, 1992.
- 2 Walsh, F. S.; Barton, C. H.; Putt, W.; Moore, S. E.; Kelsell, D.; Spurr, N.; Goodfellow, P. N. : N-cadherin gene maps to human chromosome 18 and is not linked to the E-cadherin gene. *J. Neurochem.* 55: 805-812, 1990.
- 3 Togashi, H.; Abe, K.; Mizoguchi, A.; Takaoka, K.; Chisaka, O.; Takeichi, M. : Cadherin regulates dendritic spine morphogenesis. *Neuron* 35: 77-89, 2002.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com