



Product Information Sheet

Monoclonal Anti-Neural Cell Adhesion Molecule, NCAM (Magnetic Bead conjugate)

Catalogue No. MA1068-M Immunogen

Lot No. 08A12 Growth cone enriched plasma membrane fraction from E17rat

Clone: IML-43 forebrain.

Ig type: mouse IgG1 Purification

Size: 100µg/vial Purified by the goat anti-mouse IgG affinity chromatography.

Specificity Formulation

Human, rat. 50% slurry in PBS pH 7.2 with 0.01mg NaN₃a₃ preservative.

No cross reactivity with other **Storage**

proteins. Store at 4°C for frequent use.

Recommended application Description:

Immunoprecipitation(IP) This Antagene antibody is immobilized via covalent binding of

primary amino groups to N-hydroxysuccinimide (NHS)-activated

sepharose beads. It is useful for immunoprecipitation assays.

BACKGROUND

The neural cell adhesion molecule appears on early embryonic cells and is important in the formation of cell collectives and their boundaries at sites of morphogenesis. Later in development it is found on various differentiated tissues and is a major CAM mediating adhesion among neurons and between neurons and muscle. NCAM shares many features with immunoglobulins and is considered a member of the immunoglobulin superfamily NCAM gene is mapped to 11q23.1.

REFERENCE

- 1. Cunningham, B. A.; Hemperly, J. J.; Murray, B. A.; Prediger, E. A.; Brackenbury, R.; Edelman, G. M.: Neural cell adhesion molecule: structure, immunoglobulin-like domains, cell surface modulation, and alternative RNA splicing. *Science* 236: 799-806, 1987.
- 2 D'Eustachio, P.; Owens, G. C.; Edelman, G. M.; Cunningham, B. A.: Chromosomal location of the gene encoding the neural cell adhesion molecule (N-CAM) in the mouse. *Proc. Nat. Acad. Sci.* 82: 7631-7635, 1985.
- 3. Nguyen, C.; Mattei, M. G.; Goridis, C.; Mattei, J. F.; Jordan, B. R.: Localization of the human N-CAM gene to chromosome 11 by in situ hybridization with a murine N-CAM cDNA probe. (Abstract) *Cytogenet. Cell Genet.* 40: 713 only, 1985.