



# **Product Information Sheet**

## Monoclonal Anti-Neurofilament 200 (Phos. and Nonphos.) NF200- Magnetic Bead Conjugate

Catalogue No. MA1071-M

Lot No. 08A12

Clone: NF-200

Iq type: mouse IqG1

Size: 200µl

**Specificity** 


Human, mouse, rat.

No cross reactivity with other

proteins.

**Recommended application** 

Immunoprecipitation(IP)



## **Immunogen**

C-terminal segment of enzymatically dephosphorylated pig Neurofilament 200.

### **Purification**

Purified by the goat anti-mouse IgG affinity chromatography.

#### Formulation

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN<sub>3</sub>.

## Storage

Store at 4°C for frequent use.

## Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

## **BACKGROUND**

Neurofilaments are composed of 3 neuron-specific proteins with apparent molecular masses of 68 kD (NFL), 125 kD (NFM), and 200 kD (NFH) on SDS-gel electrophoresis. Genomic clones for the largest human neurofilament protein (NF-H) were isolated, the intron/exon boundaries mapped and the entire protein-coding regions (exons) sequenced. mutations in neurofilaments have been linked to some forms of Charcot-Marie-Tooth disease (CMT)

### REFERENCE

- 1. Lees, J. F.; Shneidman, P. S.; Skuntz, S. F.; Carden, M. J.; Lazzarini, R. A.: The structure and organization of the human heavy neurofilament subunit (NF-H) and the gene encoding it. EMBO J. 7: 1947-1955, 1988.
- 2. Brownlees, J.; Ackerley, S.; Grierson, A. J.; Jacobsen, N. J. O.; Shea, K.; Anderton, B. H.; Leigh, P. N.; Shaw, C. E.; Miller, C. C.
- J.: Charcot-Marie-Tooth disease neurofilament mutations disrupt neurofilament assembly and axonal transport. Hum. Molec. Genet. 11: 2837-2844, 2002.