



Product Information Sheet

Monoclonal Anti-Phosphoserine (Sephacrose Bead Conjugate)

Catalogue No. MA1084-S

Lot No. 08A12

Clone: PS-53

Ig type: mouse IgG1

Size: 200µl

Specificity

Human, all kinds of animals.
No cross reactivity with other proteins.

Recommended application

Immunoprecipitation(IP)

Immunogen

Phosphoserine conjugated to Keyhole Limpet Hemocyanin (KLH).

Purification

Purified by the goat anti-mouse IgG affinity chromatography.

Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Storage

Store at 4°C for frequent use.

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sephacrose beads. It is useful for immunoprecipitation assays

BACKGROUND

Phosphoserine phosphatase fills an important role in the biosynthesis of serine from carbohydrates by catalyzing the last step, hydrolysis of O-phosphoserine. Phosphoserine phosphatase activity in lymphoblasts and fibroblasts was reduced to 25% of normal values. The human phosphoserine gene (PSP) is mapped to the pter-q22 region of chromosome 7.

REFERENCE

1. Koch, G. A.; Eddy, R. L.; Haley, L. L.; Byers, M. G.; McAvoy, M.; Shows, T. B. : Assignment of the human phosphoserine gene (PSP) to the pter-q22 region of chromosome 7. *Cytogenet. Cell Genet.* 35: 67-69, 1983. 2. Sparkes, R. S.; Mohandas, T.; Sparkes, M. C. : The human phosphoserine phosphatase gene (PSP) is mapped to chromosome 7 by somatic cell genetic analysis. *Cytogenet. Cell Genet.* 35: 70-71, 1983.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com