



Product Information Sheet

Monoclonal Anti-Vinculin (Magnetic Bead conjugate)

Catalogue No. MA1103-M

Immunogen

Human vinculin, purified from uterus.

Lot No. 08A12

Purification

Clone: VIN-54

Purified by the goat anti-mouse IgG affinity chromatography.

Ig type: mouse IgG1

Formulation

Size: 200µl

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN₃.

Specificity

Human, mouse, rat, chicken.

No cross reactivity with other proteins.

Storage

Store at 4°C for frequent use.

Recommended application

Immunoprecipitation(IP)

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified beads. It is useful for immunoprecipitation.

BACKGROUND

Vinculin is a cytoskeletal protein associated with the cytoplasmic face of both cell-cell and cell-extracellular matrix adherens-type junctions, where it is thought to function as one of several interacting proteins involved in anchoring F-actin to the membrane. Both human and chicken embryo sequences of vinculin contain 1,066 amino acids and, furthermore, that the 2 proteins exhibit a high level of sequence identity (greater than 95%). Vinculin is mapped to 10q22.1-q23.

REFERENCE

1. Weller, P. A.; Ogryzko, E. P.; Corben, E. B.; Zhidkova, N. I.; Patel, B.; Price, G. J.; Spurr, N. K.; Koteliensky, V. E.; Critchley, D. R. : Complete sequence of human vinculin and assignment of the gene to chromosome 10. *Proc. Nat. Acad. Sci.* 87: 5667-5671, 1990.
2. Mulligan, L. M.; Gardner, E.; Telenius, H.; Ponder, B. A. J. : Complementary physical and genetic techniques map the vinculin (VCL) gene on chromosome 10q. *Genomics* 13: 1347-1349, 1992.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com