



Polyclonal Anti-Angiopoietin-2, **ANG2 (Sephacose Bead Conjugate)**

Catalogue No. PA1005-S

Immunogen

A synthetic peptide corresponding to a sequence mapping near the C-terminal of human Angiopoietin-2 (482-496aa), identical to the related rat and mouse sequence.

Lot No. 0101012190595

Ig type: rabbit IgG

Purification

Immunogen affinity purified.

Size: 100µg/vial

Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Specificity

Human, rat.

No cross reactivity with other proteins.

Storage

Store at 4°C for frequent use.

Recommended application

Immunoprecipitation

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sephacose beads. It is useful for immunoprecipitation assays

BACKGROUND

Angiopoietin 1 and Angiopoietin 2 are important for development of the endothelium, by regulating tyrosine phosphorylation of the membrane receptor Tie 2. Angiopoietin 2 is only 60% homologous with Angiopoietin 1. Angiopoietin-2 is a naturally occurring antagonist of angiopoietin-1 that competes for binding to the TIE2 receptor and blocks ANGPT1-induced TIE2 autophosphorylation. Angiopoietin 1 binding to Tie 2 causes phosphorylation of the receptor. Angiopoietin 2 competes for this binding, and thus blocks receptor phosphorylation. Angiopoietin 2 expression occurs at sites of vascular remodelling: dorsal aorta and major aortic branches, ovary, placenta and uterus.

REFERENCE

1. Kim, I.; Kim, J.-H.; Ryu, Y. S.; Jung, S. H.; Nah, J. J.; Koh, G. Y. : Characterization and expression of a novel alternatively spliced human angiopoietin-2. *J. Biol. Chem.* 275: 18550-18556, 2000.
2. Maisonpierre, P. C.; Suri, C.; Jones, P. F.; Bartunkova, S.; Wiegand, S. J.; Radziejewski, C.; Compton, D.; McClain, J.; Aldrich, T. H.; Papadopoulos, N.; Daly, T. J.; Davis, S.; Sato, T. N.; Yancopoulos, G. D. : Angiopoietin-2, a natural antagonist for Tie2 that disrupts in vivo angiogenesis. *Science* 277: 55-60, 1997.
3. Ward, E. G.; Grosios, K.; Markham, A. F.; Jones, P. F. : Genomic structures of the human angiopoietins show polymorphism in angiopoietin-2. *Cytogenet. Cell Genet.* 94: 147-154, 2001.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com