



Product Information Sheet

Polyclonal Anti-C-C Chemokine Receptor 7, *CCR7* (Magnetic Bead Conjugate)

Catalogue No. PA1017-M

Lot No. 0101112011762

Ig type: rabbit IgG

Size: 100µg/vial

Specificity

Human, rat.

No cross reactivity with other proteins.

Recommended application

ImmunoPrecipitation (IP)

Immunogen

A synthetic peptide corresponding to a sequence mapping at the N-terminal of human CCR7, different from the related rat and mouse sequence by a single amino acid.

Purity

Immunogen affinity purified.

Contents

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN₃.

Storage

Store at 4°C for frequent use.

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified magnetic beads. It is useful for immunoprecipitation

BACKGROUND

Schweickart et al. identified CCR7 (also known as EBI1) in 1994, which is a lymphoid-specific member of the G protein-coupled receptor family. This gene, which is encoded on human chromosome 17q12-q21.2, is expressed in normal lymphoid tissues and in several B- and T-lymphocyte cell lines with 86% identity to the mouse homolog. It is also related to the receptors that recognize chemoattractants, such as interleukin-8, RANTES, C5a, and fMet-Leu-Phe. Expression of CCR7 controls homing to secondary lymphoid organs, divides human memory T cells into two functionally distinct subsets.

REFERENCE

1. Schweickart VL, Raport CJ, Godiska R, Byers MG, Eddy RL Jr, Shows TB, Gray PW. Cloning of human and mouse EBI1, a lymphoid-specific G-protein-coupled receptor encoded on human chromosome 17q12-q21.2. *Genomics*. 1994 Oct; 23(3):643-50.
2. Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature*. 1999 Oct 14; 401(6754):708-12.