



Polyclonal Anti-Cyclooxygenase-1, **COX1** (Sephacose Bead Conjugate)

Catalogue No. PA1027-S

Lot No. 05L01

Ig type: rabbit IgG

Size: 100µg/vial

Specificity

Human, mouse, rat.

No cross reactivity with other proteins.

Recommended application

(Immunoprecipitation(IP))

Immunogen

A synthetic peptide corresponding to a sequence near the N-terminal of human PTGS1(COX1), different to the related rat and mouse sequence by two amino acids.

Purification

Immunogen affinity purified.

Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Storage

Store at 4°C for frequent use.

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays

BACKGROUND

Cyclooxygenase 1(COX1), also known as Prostaglandin-endoperoxide synthase (PTGS1) or mitochondrial cytochrome c oxidase subunit 1, is the key enzyme in prostaglandin biosynthesis. The gene was approximately 40 kb long, with 11 protein-coding exons. There were 599 amino acid residues with a calculated molecular mass of approximately 68 kD. By analysis of a human/hamster somatic hybrid DNA panel, Funk et al. (1991) demonstrated that the PTGS1 gene maps to chromosome 9. Human prostaglandin endoperoxide synthase exhibited 91% amino acid identity with the sheep enzyme. Prostaglandin synthase 1 gene disruption in mice reduces arachidonic acid-induced inflammation and indomethacin-induced gastric ulceration.

REFERENCE

1. Yokoyama, C.; Tanabe, T. : Cloning of human gene encoding prostaglandin endoperoxide synthase and primary structure of the enzyme. *Biochem. Biophys. Res. Commun.* 165: 888-894, 1989. 2. Langenbach, R.; Morham, S. G.; Tiano, H. F.; Loftin, C. D.; Ghanayem, B. I.; Chulada, P. C.; Mahler, J. F.; Lee, C. A.; Goulding, E. H.; Kluckman, K. D.; Kim, H. S.; Smithies, O. : Prostaglandin synthase 1 gene disruption in mice reduces arachidonic acid-induced inflammation and indomethacin-induced gastric ulceration. *Cell* 83: 483-492, 1995.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com