



Polyclonal Anti-Tissue inhibitor of metalloproteinase 2, TIMP2 (Sepharose Bead Conjugate)

Catalogue No. PA1076-S	Immunogen
Lot No. 03A01	A peptide mapping at the C-terminal end of TIMP-2 of human origin, identical to the related rat and mouse sequence.
Ig type: rabbit	
	Purification
IgG Size: 100μg/vial	Immunogen affinity purified.
Specificity	Formulation
Human, mouse, rat.	50% slurry in PBS pH 7.2 with 0.01mg NaN ₃ a ₃ preservative.
No cross reactivity	
with other proteins.	Storage
	Store at 4°C for frequent use.
Recommended application	
Immunoprecipitation(IP)	Description:
	This Antagene antibody is immobilized via covalent binding of
	primary amino groups to N-hydroxysuccinimide
	(NHS)-activated sepharose beads. It is useful for
	immunoprecipitation assays

BACKGROUND

TIMP2 gen is encoded by 5 exons spanning 83 kb of genomic DNA. TIMP2 is 83 kilobase pairs (kb) long with exon-intron splicing sites located in preserved positions among the three members of the TIMP family. The gene for tissue inhibitor of metalloproteinases-2 is localized on human chromosome arm 17q25. TIMP-2 abrogates angiogenic factor-induced endothelial cell proliferation in vitro and angiogenesis in vivo independent of MMP inhibition.

REFERENCE

 De Clerck, Y.; Szpirer, C.; Aly, M. S.; Cassiman, J.-J.; Eeckhout, Y.; Rousseau, G. : The gene for tissue inhibitor of metalloproteinases-2 is localized on human chromosome arm 17q25. Genomics 14: 782-784, 1992. 2. Hammani, K.; Blakis, A.; Morsette, D.; Bowcock, A. M.; Schmutte, C.; Henriet, P.; DeClerck, Y. A. : Structure and characterization of the human tissue inhibitor of metalloproteinases-2 gene. J. Biol. Chem. 271: 25498-25505, 1996.
Seo, D.-W.; Li, H.; Guedez, L.; Wingfield, P. T.; Diaz, T.; Salloum, R.; Wei, B.; Stetler-Stevenson, W. G. : TIMP-2 mediated inhibition of angiogenesis: an MMP-independent mechanism. Cell 114: 171-180, 2003.