



Product Information Sheet

Polyclonal Anti-IFITM1 (*Magnetic Bead Conjugate*)

Catalogue No. PA1112-M

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminal of human IFITM1, identical to the related rat and mouse sequence.

Lot No. 01110121012126

Ig type: rabbit IgG

Purity

Immunogen affinity purified.

Size: 100µg/vial

Contents

Each vial contains 1mg/ml Magnetic Bead in PBS, pH 7.2, 0.05mg NaN₃.

Specificity

Human

No cross reactivity with other proteins.

Storage

Store at 4°C for frequent use.

Recommended application

immunoprecipitation.(IP)

Description

This Antagene antibody is immobilized by the covalent reaction of hydrazinonicotinamide-modified antibody with formylbenzamide-modified magnetic beads. It is useful for immunoprecipitation

BACKGROUND

Interferon-induced Transmembrane Protein 1 (IFITM1), also called Interferon-induced Protein 17 (IFI17). IFITM1 activity is required for primordial germ cells (PGCs) transit from the mesoderm into the endoderm, and that it appears to act via a repulsive mechanism, such that PGCs avoid Ifitm1-expressing tissues. It is mapped to Chr.11 and belongs to the family of interferon-induced transmembrane proteins (Ifitm/mil/fragilis), which encodes cell surface proteins that may modulate cell adhesion and influence cell differentiation. Interferon-inducible membrane proteins of approximately 17 kDa have been suggested to play a role in the antiproliferative activity of interferons based on their pattern of induction in interferon-sensitive and -resistant cell lines and the ability of a membrane fraction enriched in 17-kDa proteins to inhibit cell growth.

REFERENCE

1. Tanaka, S. S.; Yamaguchi, Y. L.; Tsoi, B.; Lickert, H.; Tam, P. P. L. : IFITM/Mil/Fragilis family proteins IFITM1 and IFITM3 play distinct roles in mouse primordial germ cell homing and repulsion. *Dev. Cell* 9: 745-756, 2005.
2. Deblandre, G. A.; Marinx, O. P.; Evans. S. S.; Majjaj, S.; Leo, O.; Caput, D.; Huez, G. A.; Wathélet, M. G. : Expression cloning of an interferon-inducible 17-kDa membrane protein implicated in the control of cell growth. *J. Biol. Chem.* 270: 23860-23866, 1995.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com