



Polyclonal Anti- Aquaporin 4, *AQP4* (Sepharose Bead Conjugate)

Catalogue No. PA1220-S

Lot No. 09C01

Ig type: rabbit IgG

Size: 100µg/vial

Specificity

Human, mouse, rat. No cross reactivity with other proteins.

Recommended application

(Immunoprecipitation(IP))

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminal of human AQP4, identical to the related rat and mouse sequence.

Purification

Immunogen affinity purified.

Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃a₃ preservative.

Storage

Store at 4°C for frequent use.

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays

BACKGROUND

Aquaporin 4 is found in the basolateral cell membrane of principal collecting duct cells and provide a pathway for water to exit these cells.¹ The gene of AQP4 is mapped to 18q11.2-q12.1. Similar to other aquaporins, the AQP4 gene is composed of 4 exons encoding 127, 55, 27, and 92 amino acids separated by introns of 0.8, 0.3, and 5.2 kb. Unlike other aquaporins, an alternative coding initiation sequence (designated exon 0) was located 2.7 kb upstream of exon 1. When spliced together, M1 and the subsequent 10 amino acids are encoded by exon 0; the next 11 amino acids and M23 are encoded by exon 1. AQP4 is expressed in astrocytes and is upregulated by direct insult to the central nervous system.³ AQP4 is the predominant water channel in the brain and has an important role in brain water homeostasis. It is abundant in mammalian brain and is concentrated in astrocytic foot processes at the blood-brain barrier.^{4, 5}

REFERENCE

1. Agre P, Nielsen S (1996). "The aquaporin family of water channels in kidney". *Nephrologie* 17 (7): 409–15. 2. Lu, M.; Lee, M. D.; Smith, B. L.; Jung, J. S.; Agre, P.; Verdijk, M. A. J.; Merks, G.; Rijss, J. P. L.; Deen, P. M. T. : The human AQP4 gene: definition of the locus encoding two water channel polypeptides in brain. *Proc. Nat. Acad. Sci.* 93: 10908-10912, 1996.
3. Nagelhus EA, Mathiesen TM, Ottersen OP (2004). "Aquaporin-4 in the central nervous system: cellular and subcellular distribution and coexpression with KIR4.1". *Neuroscience* 129 (4): 905–13. 4. Amiry-Moghaddam, M.; Otsuka, T.; Hurn, P. D.; Traystman, R. J.; Haug, F.-M.; Froehner, S. C.; Adams, M. E.; Neely, J. D.; Agre, P.; Ottersen, O. P.; Bhardwaj, A. : An alpha-syntrophin-dependent pool of AQP4 in astroglial end-feet confers bidirectional water flow between blood and brain. *Proc. Nat. Acad. Sci.* 100: 2106-2111, 2003.
5. Lennon, V. A.; Kryzer, T. J.; Pittock, S. J.; Verkman, A. S.; Hinson, S. R. : IgG marker of optic-spinal multiple sclerosis binds to the aquaporin-4 water channel. *J. Exp. Med.* 202: 473-477, 2005.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com