



## Polyclonal Anti- N-Cadherin (Sepharose Bead Conjugate)

**Catalogue No.** PA1328-S

**Lot No.** 013101282864

**Ig type:** rabbit IgG

**Size:** 100µg/vial

**Specificity**

Human, rat. No cross reactivity with other proteins.

**Recommended application**

(Immunoprecipitation(IP))

**Immunogen**

A synthetic peptide corresponding to a sequence at the middle region of human N-Cadherin (701-714aa), identical to the rat sequence.

**Purification**

Immunogen affinity purified.

**Formulation**

50% slurry in PBS pH 7.2 with 0.01mg NaN<sub>3</sub> preservative.

**Storage**

Store at 4°C for frequent use.

**Description:**

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays

### BACKGROUND

N-cadherin (NCAD) is a member of the cadherin cell-cell adhesion receptor family that includes P-, E-, and R-cadherin and liver cell adhesion molecule (L-CAM). N-Cadherin, also known as Cadherin-2, encodes a 907-amino acid protein that includes a 159-amino acid signal sequence. Human and mouse nucleotide sequences are 96% identical. Mouse Ncad gene consists of 16 exons dispersed over more than 200 kb of genomic DNA. Human N-cadherin gene contains 16 exons and its sequence is highly similar to both the mouse NCAD gene (including the large first and second introns) and other cadherin genes. N-cadherin is mapped to 18q11.2. Cadherin regulates dendritic spine morphogenesis.

### REFERENCE

- 1 Miyatani, S.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Takeichi, M. : Genomic structure and chromosomal mapping of the mouse N-cadherin gene. *Proc. Nat. Acad. Sci.* 89: 8443-8447, 1992.
- 2 Walsh, F. S.; Barton, C. H.; Putt, W.; Moore, S. E.; Kelsell, D.; Spurr, N.; Goodfellow, P. N. : N-cadherin gene maps to human chromosome 18 and is not linked to the E-cadherin gene. *J. Neurochem.* 55: 805-812, 1990.
- 3 Togashi, H.; Abe, K.; Mizoguchi, A.; Takaoka, K.; Chisaka, O.; Takeichi, M. : Cadherin regulates dendritic spine morphogenesis. *Neuron* 35: 77-89, 2002.

**For Research Use Only not for diagnostic and clinical use**

**Contact:** Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: [Info@antageneinc.com](mailto:Info@antageneinc.com)