



Product Information Sheet

Polyclonal Anti- L-selectin

Catalogue No. PA1359

Lot No. 01310121459124

Ig type rabbit IgG

Size 100µg/vial

Specificity

Rat, mouse

No cross reactivity with other proteins.

Recommended application

Western blot

Immunogen

A synthetic peptide corresponding to a sequence at the middle region of human L-selectin (125-141 aa), identical to the related mouse and rat sequence.

Purity

Immunogen affinity purified.

Application

	Concentration	Tested Species	Concluded Species	Antigen Retrieval
WB	1µg/ml	Rat, Ms	-	-
IHC-P	-	-	-	-
IHC-F	-	-	-	-
ICC	-	-	-	-

WB: The detection limit for L-Selectin is approximately 2.5ng/lane under non-reducing and reducing conditions.

Other applications have not been tested.

Optimal dilutions should be determined by end user.

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Reconstitution

0.2ml of distilled water will yield a concentration of 500µg/ml.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for longer time.

To reorder contact us at:

Antagene, Inc.

Toll Free: 1(866)964-2589

email: Info@antageneinc.com

BACKGROUND

L-selectin, also known as CD62L, is a cell adhesion molecule found on leukocytes. It belongs to the selectin family of proteins, which recognize sialylated carbohydrate groups. It is cleaved by ADAM17. SELL (L-selectin) is a cell surface component that is a member of a family of adhesion/homing receptors which play important roles in leukocyte-endothelial cell interactions. The molecule is composed of multiple domains: one homologous to lectins, one to epidermal growth factor, and two to the consensus repeat units found in C3/C4 binding proteins.¹ L-selectin acts as a "homing receptor" for leukocytes to enter secondary lymphoid tissues via high endothelial venules. Ligands present on endothelial cells will bind to leukocyte expressing L-selectin, slowing leukocyte trafficking through the blood, and facilitating entry into a secondary lymphoid organ at that point².

REFERENCE

- 1、"Entrez Gene: SELL selectin L (lymphocyte adhesion molecule 1)".
- 2、Cotran; Kumar, Collins (1998). Robbins Pathologic Basis of Disease. Philadelphia: W.B Saunders Company. .