



# Polyclonal Anti- Cell division control protein 42 homolog, CDC42 (Sepharose Bead Conjugate)

Catalogue No. PA1366-S

Lot No. 0131112016627

Ig type: rabbit IgG

Size: 100µg/vial

# **Specificity**

Human, rat. ,mouse, .No cross reactivity with other proteins.

# **Recommended Application**

(Immunoprecipitation(IP)

# **Immunogen**

A synthetic peptide corresponding to a sequence at the middle region of human CDC42 (121-138aa), identical to the related mouse and rat sequence.

#### **Purification**

Immunogen affinity purified.

#### Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN<sub>3</sub>a<sub>3</sub> preservative.

## Storage

Store at 4°C for frequent use.

## Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays

# **BACKGROUND**

Cell division control protein 42 homolog also known as CDC42 is a protein involved in regulation of the cell cycle. In humans, CDC42 is encoded by the CDC42 gene.CDC42 is a small GTPase of the Rho-subfamily, which regulates signaling pathways that control diverse cellular functions including cell morphology, migration, endocytosis and cell cycle progression. This protein is highly similar to Saccharomyces cerevisiae Cdc 42, and is able to complement the yeast cdc42-1 mutant. The product of oncogene Dbl was reported to specifically catalyze the dissociation of GDP from this protein. This protein could regulate actin polymerization through its direct binding to Neural Wiskott-Aldrich syndrome protein (N-WASP), which subsequently activates Arp2/3 complex. Alternative splicing of this gene results in multiple transcript variants.

### **REFERENCE**

B1. Shinjo K, Koland JG, Hart MJ, Narasimhan V, Johnson DI, Evans T, Cerione RA (December 1990). 2.Munemitsu S, Innis MA, Clark R, McCormick F, Ullrich A, Polakis P (November 1990).