



Product Information Sheet

Polyclonal Anti- Insulin-like growth factor 1, *IGF-1*

Catalogue No. PA1374

Lot No. 0131112117427

Ig type rabbit IgG

Size 100µg/vial

Specificity

Human, rat.

No cross reactivity with other proteins.

Recommended application

Western blot

Immunohistochemistry(P)

Immunogen

A synthetic peptide corresponding to a sequence at the middle region of human IGF-1 (80-96 aa), different from the mouse and rat sequence by one amino acid.

Purity

Immunogen affinity purified.

Application

	Concentration	Tested Species	Concluded Species	Antigen Retrieval
WB	1µg/ml	Hu	-	-
IHC-P	1µg/ml	Hu,Rat	Ms	By Heat
IHC-F	-	-	-	-
ICC	-	-	-	-

WB: The detection limit for IGF-1 is approximately 2.5ng/lane under non-reducing and reducing conditions.

Other applications have not been tested.

Optimal dilutions should be determined by end user.

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Reconstitution

0.2ml of distilled water will yield a concentration of 500µg/ml.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for longer time.

To reorder contact us at:

Antagene, Inc.

Toll Free: 1(866)964-2589

email: Info@antageneinc.com

BACKGROUND

Insulin-like growth factor 1 (IGF-1) also known as somatomedin C or mechano growth factor is a protein that in humans is encoded by the IGF1 gene. IGF-1 is a hormone similar in molecular structure to insulin. It plays an important role in childhood growth and continues to have anabolic effects in adults. A synthetic analog of IGF-1, mecasermin is used for the treatment of growth failure. IGF-1 consists of 70 amino acids in a single chain with three intramolecular disulfide bridges. IGF-1 has a molecular weight of 7649 daltons. Justice et al. (1990) placed the mouse IGF1 gene on chromosome 10.

REFERENCE

1. Höppener JW, de Pagter-Holthuizen P, Geurts van Kessel AH, Jansen M, Kittur SD, Antonarakis SE, Lips CJ, Sussenbach JS (1985). "The human gene encoding insulin-like growth factor I is located on chromosome 12". *Hum. Genet.* 69 (2): 157–60.
2. Jansen M, van Schaik FM, Ricker AT, Bullock B, Woods DE, Gabbay KH, Nussbaum AL, Sussenbach JS, Van den Brande JL (1983). "Sequence of cDNA encoding human insulin-like growth factor I precursor". *Nature* 306 (5943): 609–11.