



Product Information Sheet

Polyclonal Anti- Dual specificity mitogen-activated protein kinase kinase 1, *MAP2K1*

Catalogue No. PA1376

Lot No. 0131112027627

Ig type rabbit IgG

Size 100µg/vial

Specificity

Human, rat.

No cross reactivity with other proteins.

Recommended application

Western blot

Immunohistochemistry(P)

Immunocytochemistry

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminal of human MAP2K1 (353-367aa), identical to the related mouse and rat sequence.

Purity

Immunogen affinity purified.

Application

	Concentration	Tested Species	Concluded Species	Antigen Retrieval
WB	1µg/ml	Hu,Rat	-	-
IHC-P	1µg/ml	Hu,Rat	Ms	By Heat
IHC-F	-	-	-	-
ICC	1µg/ml	Hu	-	-

Other applications have not been tested.

Optimal dilutions should be determined by end user.

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Reconstitution

0.2ml of distilled water will yield a concentration of 500µg/ml.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for longer time.

To reorder contact us at:

Antagene, Inc.

Toll Free: 1(866)964-2589

email: Info@antageneinc.com

BACKGROUND

Dual specificity mitogen-activated protein kinase kinase 1 is an enzyme that in humans is encoded by the MAP2K1 gene. The protein encoded by this gene is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein kinase lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon activation by a wide variety of extra- and intracellular signals. As an essential component of the MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development. Rampoldi et al. (1997) localized the MAP2K1 gene to 15q22.1-q22.33.

REFERENCE

- 1.Rampoldi L, Zimbello R, Bortoluzzi S, Tiso N, Valle G, Lanfranchi G, Danieli GA (Mar 1998). "Chromosomal localization of four MAPK signaling cascade genes: MEK1, MEK3, MEK4 and MEKK5". Cytogenet Cell Genet 78 (3-4): 301–3.
- 2.Zheng CF, Guan KL (Jun 1993). "Cloning and characterization of two distinct human extracellular signal-regulated kinase activator kinases, MEK1 and MEK2". J Biol Chem 268 (15): 11435–9.