



Polyclonal Anti- matrix metalloproteinase-10, *MMP-10* (Sepharose Bead Conjugate)

Catalogue No. PA1380-S

Lot No. 0131112098027

Ig type: rabbit IgG

Size: 100µg/vial

Specificity

Human. No cross reactivity with other proteins.

Recommended application

(Immunoprecipitation(IP))

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminal of human MMP-10 (410-422 aa), identical to the related mouse and rat sequence.

Purification

Immunogen affinity purified.

Formulation

50% slurry in PBS pH 7.2 with 0.01mg NaN₃ preservative.

Storage

Store at 4°C for frequent use.

Description:

This Antagene antibody is immobilized via covalent binding of primary amino groups to N-hydroxysuccinimide (NHS)-activated sepharose beads. It is useful for immunoprecipitation assays

BACKGROUND

Stromelysin-2 also known as matrix metalloproteinase-10 (MMP-10) or transin-2 is an enzyme that in humans is encoded by the MMP10 gene. Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades proteoglycans and fibronectin. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.

REFERENCE

1. Muller D, Quantin B, Gesnel MC, Millon-Collard R, Abecassis J, Breathnach R (July 1988). "The collagenase gene family in humans consists of at least four members". *Biochem. J.* 253 (1): 187–92. 2. Jung JY, Warter S, Rumpler Y (1990). "Localization of stromelysin 2 gene to the q22.3-23 region of chromosome 11 by in situ hybridization". *Ann. Genet.* 33 (1): 21–3.

For Research Use Only not for diagnostic and clinical use

Contact: Antagene, Inc. | Tel: 1 (866) 964-2589 | Fax: 1 (888) 225-1868 | Email: Info@antageneinc.com