



Product Information Sheet

Peroxidase Conjugated Mouse IgG SABC Kit

Catalog No. SA1021

Size 1 kit

Product Type

Ready to use (No dilution needed)

Storage

4°C for one year. Avoid freezing.

Tested Applications

Immunohistochemical analysis of
paraffin-embedded sections, IHC(P);
Immunohistochemical analysis of
frozen-embedded sections, IHC(F);
Immunocytochemistry, IC.

Introduction

SABC (StreptAvidin-Biotin Complex) is specially designed for displaying the distribution of antigens in tissues and cells in immunochemistry and other immunodetection analyses. Streptavidin is a 47,000 dalton protein purified from the bacterium *Streptomyces avidinii*. Streptavidin has extraordinarily strong affinity to biotin molecules. The dissociation constant (K_d) of the biotin-streptavidin complex is on the order of $\sim 10^{-15}$ mol/L, a million times higher than the typical affinity between antigens and their antibodies. Streptavidin has very low non-specific binding to tissues and cells, due to its nearly neutral isoelectric point (IP=6.0~6.5). Therefore, immunohistochemical analyses based on streptavidin-biotin complex has extremely low background. Furthermore, this kit has high sensitivity because each complex it generates has a large number of peroxidase and streptavidin molecules. In brief, SABC offers high specificity, low background and ease-of-use.

Kit Components

1. 5% BSA Blocking Reagent: 12 ml, for blocking tissue sections.
2. Biotinylated Secondary Antibody (Goat Anti-mouse IgG): 12ml (10 μ g/ml). Affinity purified antibody, labeled with "long-arm" biotin (Biotinamidohexanoic acid N-hydroxysuccinimide ester, CAS# 72040-63-2).
3. SABC-POD (Peroxidase conjugated streptavidin): 12ml (20 μ g/ml). Manufactured by a proprietary method, the complex is very stable and offers superior amplification of the antigen signals.

Material Required But Not Provided

1. APES or POLY-L-LYSINE.
2. 0.02M PBS (pH 7.2~7.6): 8.5g sodium chloride, 2.8g anhydrous Na₂HPO₄ and 0.4g anhydrous NaH₂PO₄ in 1000ml of distilled water. (The weight should be adjusted accordingly if hydrous phosphates are used.)
3. 0.01 M Citrate Buffer: 3g sodium citrate dihydrate (C₆H₅Na₃O₇·2H₂O) and 0.4g citric acid monohydrate (C₆H₈O₇·H₂O) in 1000ml of distilled water.
4. DAB Chromogenic Kit (Catalog number: AR1022 or AR1025).
5. 0.1% trypsinase or the compound digest solution (Catalog number: AR0022).

To reorder contact us at:

Antagene, Inc.

Toll Free: 1(866)964-2589

Tel: (650) 964-2589

Fax: (650) 964-2519

email: Info@antageneinc.com



Product Information Sheet

Note

Mouse IgG refers to the animal origin of the primary antibody, not the origin of the specimen. This kit must be used on primary antibodies from mouse.