

# -

# **Product Information Sheet**

# FITC + POD Conjugated Human IgG SABC Kit

Catalog No. SA1084

Size 1 kit

# **Product Type**

Concentrated (Need dilution)

#### Storage

4°C for one year. Avoid freezing.

# **Tested Applications**

Immunohistochemical analysis of paraffin-embedded sections, IHC(P); Immunohistochemical analysis of frozen-embedded sections, IHC(F); Immunocytochemistry, IC.

#### **Recommended Dilution Factors**

IHC(P): 1:50-200 IHC(F): 1:50-200 IC: 1:50-200

Optimal dilutions should be determined by end users.

# Introduction

SABC (StreptAvidin-Biotin Complex) is specially designed for displaying the distribution of antigens in tissues and cells in immunochemistry and other immunodetection analyses. This kit has high sensitivity because each complex it generates has a large number of FITC or peroxidase, and streptavidin molecules. Compared to the traditional immunodetection using free FITC dyes, the SABC-FITC system greatly enhances the sensitivity and suppresses the background. FITC excitation is at 490~495nm, emission at 520~530nm.

# Kit Components

- 1. Normal rabbit serum blocking reagent: 10×, 5 ml, for blocking tissue sections.
- Biotinylated Secondary Antibody (Rabbit Anti-human IgG): 100x, 0.5ml (2mg/ml). Affinity purified antibody, labeled with "long-arm" biotin (Biotinamidohexanoic acid N-hydroxysuccinimide ester, CAS# 72040-63-2).
- SABC-FITC+Peroxidase (FITC and peroxidase conjugated streptavidin): 100x, 0.5ml (2mg/ml). Manufactured by A proprietary method, the complex is very stable and offers superior amplification of the antigen signals.
- 4. Three drop bottles (For dilution use).

### Material Required But Not Provided

- 1. APES or POLY-L-LYSINE.
- 0.02M PBS (pH 7.2~7.6): 8.5g sodium chloride, 2.8g anhydrous Na<sub>2</sub>HPO<sub>4</sub> and 0.4g anhydrous NaH<sub>2</sub>PO<sub>4</sub> in 1000ml of distilled water. (The weight should be adjusted accordingly if hydrous phosphates are used.)
- 3. 0.01 M Citrate Buffer: 3g sodium citrate dihydrate ( $C_6H_5Na_3O_7\cdot 2H_2O$ ) and 0.4g citric acid monohydrate ( $C_6H_8O_7\cdot H_2O$ ) in 1000ml of distilled water.
- 4. DAB Chromogenic Kit (Catalog number: AR1022 or AR1025).
- 5. 0.1% trypsinase or the compound digest solution (Catalog number: AR0022).

## Note

Human IgG refers to the animal origin of the primary antibody, not the origin of the specimen. This kit must be used on primary antibodies from human.

To reorder contact us at: Antagene, Inc.

Toll Free: 1(866)964-2589

Tel: (650) 964-2589 Fax: (650) 964-2519

email: Info@antageneinc.com